

Production of strange hadrons in charged jets
in Pb–Pb collisions
measured with ALICE
at the LHC

Gutachter:

Prof. Dr. Johanna Stachel

Prof. Dr. Stephanie Hansmann-Menzemer

Contents

1	Introduction	13
1.1	The standard model of particles and interactions	13
1.2	Quark-Gluon interactions	16
1.3	The Quark Gluon Plasma formation	21
1.4	Particle flow	25
1.5	Particle jets as hard probes	28
1.6	Jet quenching	30
1.7	Modification of jet observables in heavy-ion collisions	35
1.7.1	A model for jet quenching and its impact on the jet hadrochemistry	38
1.8	Identified light-flavour particle production in heavy-ion collisions	41
1.9	Analysis of identified particles in a jet-like environment	47
2	Experimental access	53
2.1	The Large Hadron Collider	53
2.2	ALICE at the LHC	55
2.3	Inner Tracking System (ITS)	56
2.4	Time Projection Chamber	59
2.5	VZERO detector	60
2.6	Zero Degree Calorimeters (ZDC)	62
2.7	Detectors for Particle Identification	62
2.8	Centrality determination	62
2.9	Event plane determination	64
2.10	Track reconstruction	65
2.11	Reconstruction of secondary vertices	67
3	Analysis of strange particles in jets in Pb-Pb collisions	69
3.1	Reconstruction of the V^0 particles	69
3.1.1	Analysis software	69
3.1.2	Data	70
3.1.3	Event selection	70
3.1.4	V^0 particle reconstruction	72
3.1.5	Signal extraction	76
3.1.6	Uncorrected p_T spectra	81
3.1.7	Analysis workflow	82

3.2	Reconstruction of jets from charged particle tracks	84
3.2.1	Some aspects of the jet reconstruction	84
3.2.2	Contribution from the Underlying Event and background fluctuations	86
3.2.3	Track selection	89
3.2.4	Combinatorial jets	89
3.2.5	Statistics after jet selection	91
3.3	Identification of V^0 particles in jet cones	92
3.4	Corrections on inclusive and intra-jet V^0 particles	94
3.4.1	V^0 reconstruction efficiency	95
3.4.2	Acceptance correction	97
3.4.3	Subtraction of particles from the Underlying Event	103
3.4.4	Subtraction of decay products (“feed-down”)	105
3.4.5	Correction of the jet transverse momenta	107
3.5	Impact of Underlying Event fluctuations on the measurements	108
3.5.1	Embedding of PYTHIA jets into PbPb events	109
3.5.2	Datasets	109
3.5.3	Jet matching	111
3.5.4	PYTHIA embedded jet spectrum	113
3.5.5	Results for the charged tracks as a crosscheck	114
3.5.6	Development of a precise V^0 background estimation derived from PYTHIA embedding	116
3.5.7	Impact of PYTHIA p_T^{jet} spectral shape on V^0 background estimation	121
3.5.8	Event-plane dependence of the V^0 background	123
3.5.9	Estimation of the systematic uncertainties	129
4	Results and discussion	143
4.1	Inclusive particles	143
4.2	$(\Lambda + \bar{\Lambda})/2K_S^0$ ratio in jets in Pb–Pb collisions	143
4.3	K_S^0 , Λ and $\bar{\Lambda}$ hadrons in “no-jet” events in Pb–Pb collisions	145
4.4	Comparison of $\Lambda + \bar{\Lambda}/2K_S^0$ ratio to other measurements	149
4.5	K_S^0 and $\Lambda + \bar{\Lambda}$ p_T spectra in jets in Pb–Pb collisions	152
5	Conclusions	157
5.1	Physics conclusions	157
5.2	Conclusions on the analysis strategy	158
	References	161
	List of Tables	173
	List of Figures	175
A	Hadronisation mechanisms	189
B	Signal extraction for inclusive V^0 particles	191
C	Signal extraction in jet cones	197

Production of strange hadrons in charged jets in Pb–Pb collisions	11
<hr/>	
D Signal extraction in the Underlying Event	207
E Technical details jet reconstruction	213
E.1 Cut selection on tracks used for jet reconstruction	213
E.2 Jet reconstruction settings	214
F Methods to estimate V^0 p_T spectrum in the Underlying Event	215
G Scaling of reference measurement in p–Pb collisions	219
H Preliminary version of the $(\Lambda+\bar{\Lambda})/2K_S^0$ ratio	223