

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b><math>SU(N)</math> gauge theory on a spatial two-dimensional twisted box</b>	<b>4</b>
<b>3</b>	<b>The torelon spectrum (non-zero electric flux)</b>	<b>7</b>
3.1	General considerations	7
3.2	Non-perturbative Results	8
3.2.1	Consistency with expectations for small and large volumes	9
3.2.2	Transition from small to large volumes	10
3.2.3	Continuity in $\tilde{\theta}$	13
3.2.4	(Non-)Existence of tachyonic instabilities	14
3.3	Derived Consequences	16
3.3.1	Conditions on the flux for the absence of phase transitions	16
3.3.2	Implications for non-commutative field theory	16
<b>4</b>	<b>The glueball spectrum (zero electric flux)</b>	<b>17</b>
4.1	General considerations	17
4.2	Non-perturbative results	18
4.2.1	$x$ -dependence of the mass gap in the $\vec{e} = 0$ sector	18
4.2.2	Filtering out the glueball from the different torelon pairs	23
4.3	Derived Consequences	25
4.3.1	$x$ -scaling and volume independence in the $\vec{e} = 0$ sector	25
4.3.2	Beating factorization for the glueball spectrum	26
<b>5</b>	<b>Conclusions</b>	<b>26</b>
<b>A</b>	<b>The Lattice simulation</b>	<b>27</b>
A.1	Lattice model	27
A.2	Determination of the spectrum	28
A.2.1	Finding plateau ranges for the mass extraction	29
A.2.2	Selection of glueball states	30
A.2.3	Suppression of finite $T$ contributions in glueball correlation functions	31
<b>B</b>	<b>Tables: non-perturbative data</b>	<b>32</b>
B.1	Lattice data for the torelon spectrum	32
B.2	Lattice data for the scalar glueball spectrum	45