

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Top-quark production at lepton colliders</b>	<b>2</b>
<b>3</b>	<b>Top-quark effective field theory</b>	<b>3</b>
3.1	Operators	4
3.2	Specific models	6
3.3	Validity	8
<b>4</b>	<b>Observables and sensitivities</b>	<b>10</b>
4.1	Cross section and forward-backward asymmetry	10
4.2	Top-quark polarization	14
4.3	CP-odd observables	17
4.4	Top-quark decay and single production	18
4.5	Bottom-quark production	21
4.6	Observables for scalar and tensor operators	23
<b>5</b>	<b>Statistically optimal observables</b>	<b>25</b>
<b>6</b>	<b>Global reach</b>	<b>30</b>
6.1	Uncertainty estimates	30
6.2	Cross section and forward-backward asymmetry	32
6.3	Statistically optimal observables	33
6.4	Global reach	35
6.5	Exploring run scenarios	37
<b>7</b>	<b>Comparison with existing limits and HL-LHC prospects</b>	<b>41</b>
7.1	Hadron-collider sensitivity to two-fermion operators	41
7.2	Hadron collider sensitivity to four-fermion operators	45
7.3	Indirect limits	46
<b>8</b>	<b>Conclusions</b>	<b>48</b>
<b>A</b>	<b>Effective-field-theory expressions for anomalous vertices</b>	<b>49</b>
<b>B</b>	<b>Additional results for the CLIC-like scenario</b>	<b>50</b>
<b>C</b>	<b>Selected results at NLO in QCD</b>	<b>53</b>
<b>D</b>	<b>Optimal observables for scalar and tensor four-fermion operators</b>	<b>53</b>
<b>E</b>	<b>Conversion to LHC TOP WG EFT conventions</b>	<b>57</b>
<b>F</b>	<b>Conversion of TopFitter limits</b>	<b>57</b>