

Patent Literature on Mosquito Repellent Inventions which Contain Plant Essential Oils – A Review

Authors

Adrian Martin Pohlit^{1,3}, Norberto Peporine Lopes³, Renata Antonaci Gama⁴, Wanderli Pedro Tadei², Valter Ferreira de Andrade Neto⁴

Affiliations

¹ Natural Products Department, National Institute for Amazon Research, Manaus, Brazil

² Health Sciences Department, National Institute for Amazon Research, Manaus, Brazil

³ Department of Chemistry and Physics, College of Pharmaceutical Sciences, São Paulo University, Ribeirão Preto, Brazil

⁴ Department of Microbiology and Parasitology, Rio Grande do Norte Federal University, Natal, Brazil

Key words

- botanical repellents
- natural products
- plant-based repellents
- repellent essential oils
- mosquito repellents

received July 2, 2010
revised Dec. 28, 2010
accepted January 3, 2011

Bibliography

DOI <http://dx.doi.org/10.1055/s-0030-1270723>
Published online February 15, 2011
Planta Med 2011; 77: 598–617
© Georg Thieme Verlag KG
Stuttgart · New York ·
ISSN 0032-0943

Correspondence

Prof. Dr. Valter Ferreira de Andrade Neto
Laboratory of Malaria and Toxoplasmosis Biology
Department of Microbiology and Parasitology
Rio Grande do Norte Federal University (UFRN)
Campus Universitário
Lagoa Nova
CEP 59072-970 Natal – RN
Brazil
Phone: + 55 84 32 15 34 37,
ext. 225, 226
Fax: + 55 84 32 11 92 10
aneto@cb.ufrn.br

Abstract

▼
Bites of mosquitoes belonging to the genera *Anopheles* Meigen, *Aedes* Meigen, *Culex* L. and *Haemagogus* L. are a general nuisance and are responsible for the transmission of important tropical diseases such as malaria, hemorrhagic dengue and yellow fevers and filariasis (elephantiasis). Plants are traditional sources of mosquito repellent essential oils (EOs), glyceridic oils and repellent and synergistic chemicals. A Chemical Abstracts search on mosquito repellent inventions containing plant-derived EOs revealed 144 active patents mostly from Asia. Chinese, Japanese and Korean language patents and those of India (in English) accounted for roughly 3/4 of all patents. Since 1998 patents on EO-containing mosquito repellent inventions have almost doubled about every 4 years. In general, these patents describe repellent compositions for use in topical agents, cosmetic products, incense, fumigants, indoor and outdoor sprays, fibers, textiles among other applications. 67 EOs and 9 glyceridic oils were individually cited in at least 2 patents. Over 1/2 of all patents named just one EO. Citronella [*Cymbopogon nardus* (L.) Rendle, *C. winterianus* Jowitt ex Bor] and eucalyptus (*Eucalyptus* L'Hér. spp.) EOs

were each cited in approximately 1/3 of all patents. Camphor [*Cinnamomum camphora* (L.) J. Presl], cinnamon (*Cinnamomum zeylanicum* Blume), clove [*Syzygium aromaticum* (L.) Merr. & L.M. Perry], geranium (*Pelargonium graveolens* L'Hér.), lavender (*Lavandula angustifolia* Mill.), lemon [*Citrus × limon* (L.) Osbeck], lemongrass [*Cymbopogon citratus* (DC.) Stapf] and peppermint (*Mentha × piperita* L.) EOs were each cited in > 10% of patents. Repellent chemicals present in EO compositions or added as pure “natural” ingredients such as geraniol, limonene, *p*-menthane-3,8-diol, nepetalactone and vanillin were described in approximately 40% of all patents. About 25% of EO-containing inventions included or were made to be used with synthetic insect control agents having mosquito repellent properties such as pyrethroids, *N,N*-diethyl-*m*-toluamide (DEET), (±)-*p*-menthane-3,8-diol (PMD) and dialkyl phthalates. Synergistic effects involving one or more EOs and synthetic and/or natural components were claimed in about 10% of all patents. Scientific literature sources provide evidence for the mosquito repellency of many of the EOs and individual chemical components found in EOs used in patented repellent inventions.

Introduction

▼ Mosquito vectors of severe tropical diseases

Severe human tropical diseases such as malaria, dengue and yellow fevers and filariasis are transmitted by the bites of infected hematophagous female mosquitoes belonging to the genera *Aedes* Meigen, *Anopheles* Meigen, *Culex* L. and *Haemagogus* L. (Diptera: Culicidae). For example, about 3.3 billion people – 1/2 of the world's population – are at risk of contracting malaria. In 2008, there were more than 247 million cases and more than

1 million deaths caused by malaria mainly in African children [1]. Human malaria is caused by infections by unicellular protozoan parasites *Plasmodium falciparum* Welch, *P. vivax* Grassi & Feletti, *P. malariae* Feletti & Grassi and *P. ovale* Stephens which are transmitted by about 20 *Anopheles* spp. Another important disease is dengue hemorrhagic fever which is a viral infection caused by several *Flavivirus* spp. (Flaviviridae) whose most important vector is *Aedes (Stegomyia) aegypti* L. Dengue and dengue hemorrhagic fevers threaten an estimated 2.5 billion people – 2/5 of the world's population – and an estimated

50 million people contract the disease per year. Around 500 000 dengue patients, most of whom are children, require hospitalization each year and around 2.5% of those affected die [2]. Another serious tropical disease which threatens about 1 billion people in 80 countries is filariasis or elephantiasis. This disease already affects an estimated 120 million people and severely incapacitates and deforms 40 million people worldwide. Filariasis is caused by infections by several roundworm species of which *Wuchereria bancrofti* Cobbold (Filariidea: Onchocercidae) is the most important and is transmitted by the bites of the common house mosquito *Cx. pipiens* L. complex, *Cx. quinquefasciatus* Say, *Aedes* and *Anopheles* spp. [3,4]. Yellow fever is an arbovirus of the *Flavivirus* genus (Flaviviridae) which is transmitted from monkeys in the jungle to humans and then from human to human by mosquitoes. The most significant yellow fever mosquito vector is *Ae. aegypti*. Despite the existence of effective vaccines, there are an estimated 200 000 cases of yellow fever and approximately 30 000 deaths attributed to this disease each year [5]. In public health initiatives which aim to limit or eradicate these and other tropical diseases, mosquito vector control methods such as repellence figure prominently among those which are employed.

Synthetic mosquito repellents DEET and alkyl phthalates

For more than 50 years, the synthetic compound DEET (*N,N*-diethyl-*m*-toluamide), has been the most effective single repellent for mosquito species and is the basis for many commercial repellent products on the market. Despite reports of severe toxic properties which can dramatically affect adults and especially young children including dermatitis, allergic reactions, neurological (seizures, coma) and cardiovascular toxicity, the risk of serious toxic effects from DEET is considered slight. Nevertheless, DEET should always be used at the lowest effective dose possible. Also, dimethyl and di-*n*-butyl phthalates (DMP and DBP, respectively), which are effective mosquito repellents and were widely used in the last century, are no longer generally recommended for use as mosquito repellents due to their toxicity [6].

Plants as sources of mosquito control agents

Plants have historically been valuable sources of agents for the control of insects [3,7]. They are the sources of the natural insecticidal and larvicidal substances nicotine (*Nicotiana* L. spp.), quassin (*Quassia amara* L.), rotenone and rotenoids (*Derris* Lour. spp. and *Lonchocarpus* Kunth spp. roots), pyrethrins like chrysanthemic acid and its derivatives present in pyrethrum [extracts of *Chrysanthemum cinerariifolium* (Trevir.) Vis. flowers] and azadirachtin (*Azadirachta indica* A. Juss. seed kernel). These and other natural insect control agents have served as the basis for the development of the structurally-related synthetic pyrethroid, nicotinoid and rotenoid classes of insecticides and piperonyl butoxide synergist. Also, pyrethroids and piperonyl butoxide synergism are the basis for a number of commercially available mosquito control products in use today [3].

Botanical repellents

Citronella essential oils (EOs) are obtained mainly from varieties of *Cymbopogon nardus* (L.) Rendle (Ceylon citronella) and *C. winterianus* Jowitt ex Bor (Java citronella). They have been used in mosquito repellency for more than a century in much of the world and are the most widely used natural repellents today [3]. Also, EOs of *Eucalyptus* L'Hér. spp. are widely used to repel insects, including mosquitoes, and contain insecticidal and repellent components *p*-menthane-3,8-diol (PMD), 1,8-cineole, α -

pinene, *p*-cymene and γ -terpinene among other active compounds [8]. According to a recent review of the scientific literature, the most frequently studied repellent EOs are those obtained from species belonging to the genera *Cymbopogon* Spreng., *Ocimum* L. and *Eucalyptus* L'Hér. spp. and a number of mosquito repellent EOs have been identified in recent years having known active repellent chemical components [7]. Mosquito repellency is believed to be due to the synergistic interactions of the chemical components in EOs. Furthermore, strong synergistic effects between EOs and isolated natural or synthetic substances have been reported [7].

The United States Environmental Protection Agency [9] has registered citronella EO, eucalyptus EO and other plant oils as safe and effective ingredients for use in topical insect repellents. However, caution is recommended in the use of EOs in general due to a number of potential toxic effects [7]. Among the important toxic effects of EOs are mutagenicity and genotoxicity. Another toxic effect is the allergenicity of EO chemical components which are controlled in the European Community and elsewhere. Interestingly, a number of repellent and insecticidal plant EOs and their chemical constituents have been evaluated using a variety of methods and are believed to be non-mutagenic [10].

A number of commercial repellent products have been developed over the past decades which utilize derivatives of plants such as EOs, fractions and their isolated chemical components and synthetic components. Thus, it is important to have a comprehensive knowledge of the commercially significant uses of EOs in mosquito repellent inventions and to the best of our knowledge the patent literature on this topic has not been reviewed. The aim of the present review is to explore and analyze patent literature on mosquito repellent inventions which make use of or are based wholly on plant EOs and/or their chemical components. A secondary aim is to analyze the scientific bases and relevancy of the use of plant EOs and chemical components of these oils in patented mosquito repellent formulations.

Chemical Abstracts Search Criteria



A Chemical Abstracts search of the patent literature for the period of 1991 through May, 2010 was performed using SciFinder Scholar® [11]. Combinations of the key words "repellent", "mosquito" and "essential oil" generated an initial set of approximately 160 patents. This set was further refined. Patents describing mosquito-tocidal/larvicidal inventions, but having no stated claim or use as mosquito repellents were eliminated. Also, patented mosquito repellent inventions which did not make use of at least one commercially-obtained, plant-derived oil [a volatile (essential) oil, a concrete, or a pressed oil] or which did not describe the preparation of an EO by physical means for use within the patent were eliminated. This approach eliminated patents presenting solvent extraction performed on mixtures of several plant materials followed, for example, by distillation and evaporation as a means of obtaining the mosquito repellent invention. Also, by this criterion, incense and other smoke-generating inventions made from mosquito repellent plants but having no (added) EO in the composition were eliminated. Applying the above criteria led to a data set of 144 patents describing plant oil-containing mosquito repellent inventions which was the basis for the analysis presented below.




Fig. 1 Analysis of EO-containing patented mosquito repellent inventions by patenting country, patent language, patent type and utility.

Patenting Countries and Patent Families: Trends

The contribution of China, Japan, Korea and India to the overall number of patented EO-containing mosquito repellent inventions is significant. These countries accounted for, respectively, 37, 15, 13 and 8% (73%) of all patents (◉ Fig. 1). An interesting trend is that all 53 Chinese, 17 of 18 Korean, 20 of 22 Japanese and 9 of 11 Indian patents have only been deposited in their countries of origin and in general have not been followed-up by later patents. Also, all Belgian, Brazilian, German and Polish patents followed the rule of single deposits without follow-up patents. In contrast, industrial, academic and other patents originating in Australia, Canada, the United Kingdom and the United States gave rise to larger patent families, made greater use of the World Patent System and led to deposits of patents on EO-containing repellents in multiple countries as a rule. Finally, more than 75% of all patented repellent inventions are indexed as agrochemical bioregulators and over 80% describe chemical compositions (often together with preparation details) (◉ Fig. 1).

EOs Used in Patented Mosquito Repellent Inventions

Plant taxonomic information [12], extraction methods, chemical composition, literature sources [8,10,13–115] and frequency of use for 67 EOs and 9 glyceridic oils which were individually cited in at least 2 (1.4%) of the total number of patents are presented in ◉ Table 1. Approximately 60% of all patents named just one plant EO. One patent claimed to use up to 44 plant oils in formulations. Citronella (34.7%) and eucalyptus (30.6%) EOs were each named in about 1/3 of all patented repellent inventions followed by (species, % of patents): lavender (*Lavandula angustifolia* Mill., 21.5%), peppermint (*Mentha × piperita* L., 16.7%), clove [*Syzygium aromaticum* (L.) Merr. & L.M. Perry, 15.3%], lemongrass [*Cymbopogon citratus* (DC.) Stapf, 14.6%], cinnamon (*Cinnamomum zeylanicum* Blume, 12.5%), geranium (*Pelargonium graveolens* L'Hér., 11.8%), camphor [*Cinnamomum camphora* (L.) J. Presl, 11.1%] and lemon [*Citrus × limon* (L.) Osbeck, 11.1%] EOs. A number of EOs were cited in only one patented repellent invention (and were not included in ◉ Table 1). Given the importance

Table 1 Botanic information, extraction methods, major chemical components, literature sources and frequency (%) of use of plant essential and glyceridic oils in patented mosquito repellent inventions, 1991 – May, 2010.

Oil	Species	Family	Part	Method	Major components	Lit.	% Patents
Essential Oils							
Ambrette	<i>Abelmoschus moschatus</i> Medik.	Malvaceae	whole seed	steam distilled	<i>Z</i> E, <i>6</i> E-farnesyl acetate, <i>Z</i> -7-hexadecen-16-olide, β -farnesene	[13, 14]	1.4
Angelica	<i>Angelica archangelica</i> L.	Apiaceae	root	steam distilled	ligustilide, α & β -pinene, carvacrol , 3-carene , limonene , β -phellandrene, 15-pentadecanolide	[15, 16]	1.4
Anise	<i>Pimpinella anisum</i> L.	Apiaceae	fruit, seed	steam distilled	<i>trans</i> -anethole	[17]	4.2
Artemisia	<i>Artemisia argyi</i> H. Lévl. & Vaniot	Asteraceae	leaf	steam distilled	germacrene D, α -phellandrene, α-myrcene , 1,8-cineole , borneol , terpinol, spathulenol	[18, 19]	4.9
Basil	<i>Ocimum basilicum</i> L.	Lamiaceae	leaf, flower top	steam distilled	estragole, limonene , fenchone , linalool , eugenol <i>E</i> -methyl cinnamate, 1,8-cineole	[20–22]	4.9
Bay laurel	<i>Laurus nobilis</i> L.	Lauraceae	leaf	steam distilled	1,8-cineole , sabinene, α -terpinyl acetate, linalool	[23, 24]	1.4
Bergamot	<i>Citrus × bergamia</i> Risso	Rutaceae	fresh or dried peel	pressed	limonene , linalyl acetate, β-pinene , γ-terpinene , linalool	[24]	1.4
Camphor	<i>Cinnamomum camphora</i> (L.) J. Presl	Lauraceae	wood, bark, leaf	steam distilled	1,8-cineole , α-terpineol , α-pinene , linalool , camphor , sabinene	[10]	11.1
Cassia	<i>Cinnamomum cassia</i> (L.) C. Presl	Lauraceae	leaf, bark, stalk	steam distilled	<i>bark</i> : <i>E</i>-cinnamaldehyde , methyl <i>o</i>-salicylate ; <i>leaf</i> : 3-methoxy-1,2-propanediol, <i>E</i>-cinnamaldehyde , <i>o</i> -methoxy-cinnamaldehyde	[25, 26]	2.1
Catnip, catmint	<i>Nepeta cataria</i> L.	Lamiaceae	dry leaf, stem	steam distilled	nepetalactone , 1,8-cineole , α -humulene, α -pinene, <i>E</i>-geraniol , β -caryophyllene, citronellol	[20, 27, 28]	4.9
Cedar	<i>Cedrus</i> Trew (<i>Cupressus</i> L., <i>Juniperus</i> L.) spp.	Pinaceae (Cupressaceae)	wood	steam distilled	thujopsene, eudesmol, <i>E</i> (+)- α -atlantone; α , β & γ -himachalenes; α - & β -cedrenes; limonene , β -phellandrene, α & β-pinene , 3-carene ; <i>p</i> -methyl- Δ -3-tetrahydro & <i>p</i> -methyl acetophenones; hinokitiol, carvacrol	[29]	9.7
Chamomile	<i>Chamaemelum nobile</i> (L.) All. <i>Matricaria recutita</i> L.	Asteraceae	seed, leaf, flower	steam distilled	<i>Roman</i> : isobutyl, isoamyl & 2-methylpentyl angelates, α-pinene <i>German</i> : <i>E</i> - β -farnesene, <i>E</i> , <i>E</i> - α -farnesene, α-bisabolol , α -bisabolol oxides A & B	[20, 28, 30]	2.1
Chrysanthemum	<i>Chrysanthemum indicum</i> L.	Asteraceae	dry flower	steam distilled	verbenol , 2-(2,4-hexadiynylidene)-1,6-dioxaspiro[4.4]non-3-ene, 1,8-cineole , α-pinene , camphor , borneol , bornyl acetate	[31, 32]	2.8
Cinnamon	<i>Cinnamomum zeylanicum</i> Blume	Lauraceae	bark, leaf	steam distilled	eugenol , cinnamaldehyde	[25, 33, 34]	12.5
Citronella	<i>Cymbopogon nardus</i> (L.) Rendle, <i>C. winterianus</i> Jowitt ex Bor	Poaceae	leaf	steam distilled	citronellal , geraniol , citronellol , geranyl acetate	[35, 36]	34.7
Citrus	<i>Citrus × limon</i> (L.) Osbeck, <i>C. × sinensis</i> (L.) Osbeck, <i>C. × aurantifolia</i> L.	Rutaceae	peel	pressed, steam distilled	see orange, lemon, lime oil compositions	–	1.4
Clove	<i>Syzygium aromaticum</i> (L.) Merr. & L. M. Perry	Myrtaceae	flower bud	steam distilled	eugenol , caryophyllene, eugenyl acetate	[37]	15.3
Coriander	<i>Coriandrum sativum</i> L.	Apiaceae	fruit, seed	steam distilled	linalool , geraniol , geranyl acetate , 2-decenal, 3-dodecenal	[20, 38, 39]	2.1
Cypress	<i>Cupressus sempervirens</i> L.	Cupressaceae	needle, twig	steam distilled	sabinene, α-pinene , terpinen-4-ol , limonene	[40, 41]	4.2
Dill	<i>Anethum graveolens</i> L.	Apiaceae	seed, leaf, stem	steam distilled	carvone , limonene , α -phellandrene, α-pinene , <i>cis</i> -dihydrocarvone	[42, 43]	1.4
Eucalyptus	<i>Eucalyptus</i> L'Hér. spp.	Myrtaceae	leaf	steam distilled	1,8-cineole , <i>p</i>-menthane-3,8-diol , α-pinene , <i>p</i>-cymene , γ-terpinene , eucamalol, <i>allo</i> -ocimene, citronellol , α-terpineol	[8]	30.6
Fennel	<i>Foeniculum vulgare</i> Mill.	Apiaceae	fruit	steam distilled	<i>E</i> -anethole, (+)-fenchone , α -phellandrene, (±)-limonene , estragole	[24]	4.9
Garlic	<i>Allium sativum</i> L.	Amaryllidaceae	bulb	steam distilled	diallyl disulfide, diallyl trisulfide, methyl allyl trisulfide	[44, 45]	6.3
Geranium	<i>Pelargonium graveolens</i> L'Hér.	Geraniaceae	leaf, stem	steam distilled	2-phenylethanol, geraniol , citronellol , geranyl acetate	[46]	11.8

continued next page

Table 1 Botanic information, extraction methods, major chemical components, literature sources and frequency (%) of use of plant essential and glyceric oils in patented mosquito repellent inventions, 1991 – May, 2010. (continued)

Oil	Species	Family	Part	Method	Major components	Lit.	% Patents
Essential Oils							
Ginger	<i>Zingiber officinale</i> Roscoe	Zingiberaceae	rhizome	steam distilled	geranial , α -zingiberene, <i>E,E</i> - α -farnesene, neral, ar-curcumene, geraniol	[47, 48]	1.4
Grapefruit	<i>Citrus reticulata</i> Blanco	Rutaceae	peel	pressed	limonene , geranial , neral	[49]	2.1
Guaiaac wood	<i>Bulnesia sarmientoi</i> Lorentz ex Griseb.	Zygophyllaceae	wood	steam distilled	bulnesol, guaial, α -bulnesene	[50]	1.4
Hiba	<i>Thujopsis dolabrata</i> (Thunb. ex L. f.) Siebold & Zucc.	Cupressaceae	wood	steam distilled	sabinene, 4-terpineol, thujopsene, hinokitiol, α -thujaplicine, carvacrol	[51–53]	2.1
Ho leaf	<i>Cinnamomum camphora</i> (L.) J. Presl	Lauraceae	leaf	steam distilled	1,8-cineole , α-terpineol , linalool , camphor , safrole, sabinene, nerolidol	[10, 54, 55]	1.4
Hyssop	<i>Hyssopus officinalis</i> L.	Lamiaceae	leaf, flower	steam distilled	sabinene, pinocamphene, isopinocamphene, isopinocamphone, pinocarvone, <i>cis</i> & <i>trans</i> -pinocamphones, β-pinene , 1,8-cineole , camphor , linalool	[56–58]	1.4
Jasmine	<i>Jasminum officinale</i> L.	Oleaceae	flower	solvent extracted	linalool , benzyl acetate, methyl & benzyl benzoates, methyl anthranilate, Zjasmone, eugenol	[59, 60]	3.5
Juniper	<i>Juniperus communis</i> L.	Cupressaceae	fruit	steam distilled	α-pinene , myrcene , sabinene, germacrene D	[61]	1.4
Lady-of-the-night	<i>Cestrum nocturnum</i> L.	Solanaceae	flower	solvent extracted, steam distilled	phenylethyl alcohol, benzyl alcohol, eugenol	[62]	2.1
Lavender	<i>Lavandula angustifolia</i> Mill.	Lamiaceae	flower	steam distilled	linalool , linalyl acetate, lavandulyl acetate, α-terpineol , geranyl acetate , terpinen-4-ol , 1,8-cineole	[63, 64]	21.5
Lemon	<i>Citrus × limon</i> (L.) Osbeck	Rutaceae	peel	pressed	limonene, β -pinene, γ -terpinene	[65]	11.1
Lemon eucalyptus	<i>Eucalyptus citriodora</i> Hook.	Myrtaceae	leaf, twig	steam distilled	citronellal , citronellol	[8, 66]	8.3
Lemongrass	<i>Cymbopogon citratus</i> (DC.) Stapf	Poaceae	leaf	steam distilled	geranial , neral, myrcene	[41]	14.6
Lemon tea tree	<i>Leptospermum petersonii</i> F. M. Bailey	Myrtaceae	leaf	steam distilled	neral, geranial , γ-terpinene , geraniol , geranyl acetate , α-pinene , citronellal , terpinolene	[67]	1.4
Lime	<i>Citrus × aurantifolia</i> L.	Rutaceae	peel	steam distilled, pressed	<i>D</i> -dihydrocarvone, <i>D</i>-limonene , α-terpineol	[68]	2.1
Manuka	<i>Leptospermum scoparium</i> Forst. & Forst.	Myrtaceae	leaf, stem	steam distilled	leptospermone, <i>trans</i> -calamenene, flavesone, 1,8-cineole , α-pinene	[69, 70]	1.4
Marjoram	<i>Origanum majorana</i> L.	Lamiaceae	leaf, flower	steam distilled	<i>p</i>-cymene , γ-terpinene , terpinen-4-ol , linalool , <i>cis</i> -sabinene hydrate	[71, 72]	2.1
May chang/ Litsea	<i>Litsea cubeba</i> (Lour.) Pers.	Lauraceae	fruit	steam distilled	neral, <i>R</i>-(+)-limonene , geranial , citronellal	[46, 73, 74]	4.9
Melaleuca/ Tea tree	<i>Melaleuca alternifolia</i> Cheel	Myrtaceae	leaf	steam distilled	terpinen-4-ol , γ-terpinene , α-terpinene , 1,8-cineole	[75]	9.7
Mint, mentha	<i>Mentha</i> L. spp.	Lamiaceae	leaf, flower	steam distilled	menthone , menthol , 1,8-cineole , 4-terpineol	[76]	9.7
Orange	<i>Citrus × sinensis</i> (L.) Osbeck	Rutaceae	peel	pressed	limonene , myrcene	[77]	4.2
Palmarosa	<i>Cymbopogon martini</i> (Roxb.) Will. Watson	Poaceae	dry leaf	steam distilled	geraniol , geranyl acetate , geranial	[78]	4.9
Parsley (Curl leaf)	<i>Petroselinum crispum</i> (Mill.) Fuss	Apiaceae	leaf, stem, seed	steam distilled	β -phellandrene, myristicin, α & β-pinene , myrcene	[79]	1.4
Patchouli	<i>Pogostemon cablin</i> (Blanco) Benth.	Lamiaceae	dry, fermented leaf	steam distilled	(-)-patchoulol, α -guaiene, seychellene, β -caryophyllene, α & β -patchoulenes, selinene, α -bulnesene, norpatchoulol, pogostol	[80]	4.2
Pepper	<i>Piper nigrum</i> L.	Piperaceae	fruit	steam distilled	<i>E</i> - β -caryophyllene, limonene , β-pinene	[81, 82]	2.1
Peppermint	<i>Mentha × piperita</i> L.	Lamiaceae	aerial part	steam distilled	isomenthol, <i>p</i>-menthone , isomenthyl & menthyl acetates	[83, 84]	16.7
Pine	<i>Pinus sylvestris</i> L.	Pinaceae	twig, bud	steam distilled	3-carene , α & β-pinene , α -cadinol, camphene	[85, 86]	8.3
Rose	<i>Rosa × damascena</i> Mill., <i>R. × centifolia</i> L.	Rosaceae	petal	steam distilled, solvent extracted	2-phenethyl alcohol, citronellol , geraniol , linalool , nonadecane	[30, 87]	6.3

continued next page

Table 1 Botanic information, extraction methods, major chemical components, literature sources and frequency (%) of use of plant essential and glyceridic oils in patented mosquito repellent inventions, 1991 – May, 2010. (continued)

Oil	Species	Family	Part	Method	Major components	Lit.	% Patents
Essential Oils							
Rosemary	<i>Rosmarinus officinalis</i> L.	Lamiaceae	flower	steam distilled	verbenone, camphor, borneol, bornyl acetate, α-terpineol, terpinen-4-ol	[41, 88]	9.0
Salvia (Sage)	<i>Salvia</i> L. spp.	Lamiaceae	leaf, flower, aerial part	steam distilled	camphor, 1,8-cineole, α-pinene, camphene, limonene, linalool, myrcene, β-caryophyllene, caryophyllene oxide, spathulenol, viridiflorol, 3-carene, α-bisabolol, geraniol, linalyl acetate, γ-muurolene, α-thujone, manool	[89–92]	1.4
Sandalwood	<i>Santalum</i> L. spp.	Santalaceae	heartwood	steam distilled	α, β & ϵ-<i>epi</i>-santalenes; α, β & <i>epi</i>-β-santalals; α-santalol; spiro, β, <i>E</i>-β & <i>epi</i>-β santalols, <i>trans</i>-α-bergamotene, <i>trans</i>-α-bergamotol	[80, 93]	3.5
Sour (bitter) orange	<i>Citrus \times aurantium</i> L.	Rutaceae	peel	pressed	limonene	[94]	3.5
Spearmint	<i>Mentha spicata</i> L.	Lamiaceae	flower	steam distilled	carvone, limonene	[95]	2.8
Star anise	<i>Illicium verum</i> Hook. f.	Schisandraceae	seed	steam distilled	<i>E</i> -anethole, 4-allylanisole	[96, 97]	see anise*
Tagetes	<i>Tagetes minuta</i> L.	Asteraceae	leaf, stalk, flower	steam distilled, solvent extracted	<i>E</i> & <i>Z</i> - β -ocimenes, limonene, α-terpineol , dihydrotagetone, <i>Z</i> -tagetone, <i>E</i> & <i>Z</i> -tagetenes	[98]	2.1
Tarragon	<i>Artemisia dracunculul</i> L.	Asteraceae	leaf, flower	steam distilled	sabinene, elemicine, methyl eugenol	[99]	1.4
Thyme	<i>Thymus vulgaris</i> L.	Lamiaceae	flower, leaf	steam distilled	<i>p</i>-cymene, geraniol, <i>cis</i>-thujone, thymol, carvacrol	[41]	7.6
Turmeric	<i>Curcuma longa</i> L.	Zingiberaceae	rhizome	steam distilled, solvent extracted	α -phellandrene, 1,8-cineole, terpinolene , zingiberene, β -sesquiphellandrene, α & β -turmerones	[41]	2.8
Verbena	<i>Lippia triphylla</i> (L'Hér.) Kuntze	Verbenaceae	leaf	steam distilled	geraniol, neral, limonene, geraniol	[100]	1.4
Vetiver	<i>Vetiveria zizanioides</i> (L.) Nash	Poaceae	root	steam distilled	khusimol, isonootkatool, β -vetivenene, α & β -vetivones	[101]	2.1
Violet	<i>Viola odorata</i> L.	Violaceae	leaf, flower	solvent extracted	nona-2,6-dienal, <i>cis</i> -hex-3-en-1-ol, hexadec-1-ene, pentadec-3-enal, octadec-1-ene, hexadecanoic & octadeca-9,12-dienoic acids	[102]	1.4
Wintergreen	<i>Gaultheria procumbens</i> L.	Ericaceae	leaf	steam distilled	methyl salicylate	[103]	6.3
Ylang-Ylang	<i>Cananga odorata</i> (Lam.) Hook. f. & Thomson	Annonaceae	flower	steam distilled	linalool , benzyl acetate, benzyl benzoate, benzyl salicylate	[41]	2.8
Glyceridic Oils							
Castor	<i>Ricinus communis</i> L.	Euphorbiaceae	seed	pressed	ricinolic, linoleic, oleic acids	[104]	4.9
Evening primrose	<i>Oenothera biennis</i> L.	Onagraceae	seed	pressed	linalool ; palmitic, stearic, oleic , linoleic & γ -linolenic acids	[105, 106]	1.4
Ligusticum/ Chuanxiong	<i>Ligusticum chuanxiong</i> Hortorum ex Qiu, et al. in Qiu	Apiaceae	root	CO ₂ extracted	butylphthalide, 2-propenyl phenoxyacetate, 3-isobutyliden phthalide, palmitic & octadecenoic acids	[107]	1.4
Mustard	<i>Brassica</i> L. spp.	Brassicaceae	seed	Pressed	erucic, oleic , linoleic, linolenic, palmitic & stearic acids	[108]	2.1
Neem	<i>Azadirachta indica</i> A. Juss.	Meliaceae	seed	pressed	salanin, nimbinin, nimbin, nimbidiol, azadirachtin; palmitic, stearic, oleic , linoleic acids	[104, 109]	8.3
Olive	<i>Olea europaea</i> L.	Oleaceae	fruit	pressed	oleic , palmitic, linoleic & stearic acids	[110]	2.1
Perilla	<i>Perilla frutescens</i> (L.) Britton	Lamiaceae	seed	pressed	oleic , linoleic & linolenic acids, S-limonene , perillaldehyde , elsholzia- ketone, naginataketone, perillaketone, myristicin, dillapiol, elemicin, isoeogonaketone, perillene, egomaketone, shisofuran	[111–113]	1.4
Sesame	<i>Sesamum indicum</i> L.	Pedaliaceae	seed	pressed	linoleic, oleic , palmitic & stearic acids	[114]	3.5
Soybean	<i>Glycine max</i> (L.) Merr.	Fabaceae	seed	solvent extracted	linoleic, oleic , palmitic, α -linolenic & stearic acids	[115]	4.2

Note: chemical components in **bold text** have proven mosquito repellent and deterrent properties according to literature and patent sources which are summarized in **Table 4**. Tropicos, botanical information system at the Missouri Botanical Garden [12] was used as a general reference on taxonomic information in this table

of potentially novel sources of mosquito repellent botanicals it is important to mention the following EOs: abies [*Abies spectabilis* (D. Don) Spach], almond [*Prunus dulcis* (Mill.) D.A. Webb], Austr-

lian yuzu (*Citrus junos* Siebold ex Tanaka), black pine (*Pinus nigra* Arnold), *Blumea lacera* (Burm. f.) DC., calamus (*Acorus calamus* L.), Canada fleabane [*Conyza canadensis* (L.) Cronquist], carda-

Table 2 Uses and applications, EOs and other components of patented mosquito repellent inventions which claimed synergist effects.

Use or technology	Plant EO	Plant species	Non-EO component	Source
Applied films	–	<i>Backhousia citriodora</i> F. Muell. <i>Melaleuca ericifolia</i> Sm.	–	[116]
Coatings	Wintergreen Camphor	<i>Gaultheria procumbens</i> L. <i>Cinnamomum camphora</i> (L.) J. Presl	pyrethroids	[117]
Floor wash	Eucalyptus Lemon eucalyptus Citronella	<i>Eucalyptus</i> L'Hér. spp. <i>Eucalyptus citriodora</i> Hook. <i>Cymbopogon winterianus</i> Jowitt ex Bor	–	[118]
Incense	Eucalyptus Cassia	<i>Eucalyptus</i> L'Hér. spp. <i>Cinnamomum cassia</i> (L.) C. Presl	pyrethroids	[119]
Microcapsules for textiles	Eucalyptus	<i>Eucalyptus</i> L'Hér. spp.	DEET, isobornyl derivatives	[120]
Not specified	Rose Jasmine Limonene	<i>Rosa</i> L. sp. <i>Jasminum officinale</i> L. synergist natural product	natural pyrethrins	[121]
Spray/Fumigation on grass, shrubs, trees	Citronella Neem Eucalyptus	<i>Cymbopogon</i> Spreng. sp. <i>Azadirachta indica</i> A. Juss. <i>Eucalyptus</i> L'Hér. spp.	dimethyl phthalate, allyl sulfate	[122]
Vaporizer/Fumigant	Lippia Geranium Lemon eucalyptus Basil etc.	<i>Lippia</i> L. sp.+: <i>Pelargonium graveolens</i> L'Hér. <i>Eucalyptus citriodora</i> Hook. <i>Ocimum basilicum</i> L.	–	[123]

mom [*Elettaria cardamomum* (L.) Maton], cork tree (*Phellodendron amurense* Rupr.), *Elsholtzia hunanensis* Hand.-Mazz., flos loniceræ japonicæ or jin yin hua (*Lonicera japonica* Thunb. ex Murray flower), fructus forsythiæ [*Forsythia suspensa* (Thunb.) Vahl], galbanum (*Ferula galbaniflua* Boiss. & Buhse), herba schizonepetæ [*Schizonepeta* (Benth.) Briq. sp.], hibiscus (*Hibiscus* L. sp.), larch (*Larix* Mill. sp.), lemon balm (*Melissa officinalis* L.), lovage (*Levisticum officinale* W.D.J. Koch), lilac (*Syringa vulgaris* L.), *Limnanthes alba* Hartw. ex Benth. seed, linaloe wood (*Bursera delpechiana* Poiss. ex Engl.), *Michelia* × *alba* DC. leaf, myrtle (*Myrtus communis* L.), *Ocimum canum* Sims, onion (*Allium cepa* L.), oregano (*Origanum vulgare* L.), peach [*Prunus persica* (L.) Batsch], pennyroyal (*Mentha pulegium* L.), pepper (*Capsicum annum* L. var. *annuum*), petitgrain (*Citrus aurantium* L. var. *amara*), pimento [*Pimenta dioica* (L.) Merr.], pine needle (*Pinus* L. sp.), *Rhodomyrtus tomentosa* (Aiton) Hassk., rue (*Ruta graveolens* L.), *Stephania sinica* Diels, tansy (*Tanacetum* L. sp.), *Torreya grandis* Fortune ex Lindl. and valerian (*Valeriana* L. sp.). Nevertheless, we will not systematically go into details of the composition and repellency of these EOs in this review.

Generally, patent literature on mosquito repellent inventions treats fragrant extracts as EOs whether these extracts are produced using an initial solvent extraction step (such as in the production of concretes), by steam distillation, hydrodistillation, direct distillation or pressing (resulting in proper EOs). Herein, the term EO refers broadly to fragrant extracts obtained using these different extraction techniques. Importantly, fragrant oils and partially volatile balsamic oils which have volatile and non-volatile (glyceridic) chemical components in their compositions (Table 1) and mosquito repellent properties, such as evening primrose (*Oenothera biennis* L.), perilla [*Perilla frutescens* (L.) Britton], ligusticum (*Ligusticum chuanxiong* Hortorum ex Qiu, et al. in Qiu) and copaiba (*Copaifera* L. spp.) oils are used in mosquito repellent patented formulations together with plant EOs.

Glyceridic Oils Used in Patented Mosquito Repellent Inventions

Importantly, plant glyceridic oils, several of which are known to possess mosquito repellency, were used in patented formulations as carriers or active ingredients which were associated with prolonged repellent action. The most used glyceridic oils in patented EO-containing repellent inventions were neem/margarosa (*Azadirachta indica* A. Juss., 8.3%), castor (*Ricinus communis* L., 4.9%), soybean [*Glycine max* (L.) Merr., 4.2%] and sesame (*Sesamum indicum* L., 3.5%) oils (Table 1).

Synergist Effects Associated with EO-Containing Compositions

Synergist interactions were claimed to be operating in 8% of EO-containing patents (Table 2) [116–123]. For example, EO of a species of *Lippia* L. and any one of a number of EOs formulated into a slowly evaporating hydrocarbon soluble composition was said to modify neuronal activity in invertebrates such as adult mosquitoes and produce repellent activity comparable to commercial pyrethroids [123]. Another example of a synergist formulation involved almost equal amounts of eucalyptus (*Eucalyptus* spp.) and cassia [*Cinnamomum cassia* (L.) C. Presl] EOs together with an emulsifying agent and butyl acetate solvent which was meant to be used with pyrethroids in incense formulations [119]. Still another example of a synergist repellent meant for use with pyrethroids (allethrin, dimefluthrin) used EOs of wintergreen (*Gaultheria procumbens* L.) 30–50 (w/w) and camphor (*Cinnamomum camphora* (L.) J. Presl) 10–35 (w/w), emulsifier and solvent [117] (Table 2).

Scientific Basis for Mosquito Repellency of EOs in Patented Inventions

▼
The scientific literature on the mosquito repellency of EOs used in patents provides important insights (► **Table 3**) [8, 37, 41, 46, 61, 68, 73, 85, 88, 93, 124–177]. Firstly, citronella (*Cymbopogon nardus*, *C. winterianus*), eucalyptus (*Eucalyptus* spp.) and lemon eucalyptus (*E. citriodora* Hook.) EOs which were cited in many patented inventions have been the subject of a number of studies in which repellency against species of *Culex*, *Anopheles* and *Aedes* in some cases comparable to DEET have been reported for these oils alone or in formulations [138, 140, 143, 178]. Furthermore, many EOs used in patented inventions have quite significant repellent properties according to published studies such as: bay laurel (*Laurus nobilis* L.) [137], camphor (*Cinnamomum camphora*) [132], cassia (*Cinnamomum cassia*) [132] EOs against *Ae. aegypti*; lemon (*Citrus × limon*) EO [128, 152] against *An. stephensi* Liston, catnip (*Nepeta cataria* L.) EO [128], lemongrass (*Cymbopogon citratus*) [128], may chong/litsea [*Litsea cubeba* (Lour.) Pers.] [128, 179], tagetes (*Tagetes minuta* L.) [128], violet (*Viola odorata* L.) EOs [128] against *Ae. aegypti*, *An. stephensi* and *Cx. quinquefasciatus*; peppermint (*Mentha × piperita*) EO on human skin against *An. annularis* van der Wulp, *An. culicifacies* Giles and *Cx. quinquefasciatus* [160]; sandalwood (*Santalum* L. spp.) EO formulations against a *Culex* sp. [177]; geranium (*Pelargonium graveolens*) oil formulations [137, 144, 147, 148] against species of *Culex*, *Anopheles* and *Aedes*; thyme (*Thymus vulgaris* L.) EO against *Cx. quinquefasciatus* [128, 137, 162]; marjoram (*Origanum majorana* L.) and juniper (*Juniperus communis* L.) EOs against *Cx. pipiens pallens* Coquillett [151]; and wintergreen (*Gaultheria procumbens*) EO against species of *Culex* and *Aedes* [131].

A broad-scale screening of plant oils against *Ae. aegypti*, *An. stephensi* and *Cx. quinquefasciatus* evaluated protection periods and percent of repellence on human skin as compared to 20% DEET [128]. In general, *Ae. aegypti* was the most difficult species to repel, followed by *An. stephensi* and finally *Cx. quinquefasciatus* for both oils and controls. The control (DEET) exhibited a protection period (PP) of 6 h and percent repellency (R%) of 46% against *Ae. aegypti*, whereas against *An. stephensi* and *Cx. quinquefasciatus* protection was for 8 h at 100% repellency [128] (► **Table 3**). While the most active EOs against all three mosquito species were cited above, the following oils were active against *An. stephensi* and *Cx. quinquefasciatus*, but not significantly active against *Ae. aegypti*: chamomile (*Chamaemelum nobile*), cinnamon (*Cinnamomum verum*), galbanum (*Ferula galbaniflua*), jasmine (*Jasminum grandiflorum*), lavender (*Lavandula angustifolia*), pepper (*Piper nigrum*), rosemary (*Rosmarinus officinalis*), sandalwood (*Santalum album*) and soybean (*Glycine max*) [128]. Another group of oils in this study actively repelled only *Cx. quinquefasciatus*: cedar (*Cedrus*, *Cupressus* and *Juniperus* spp.), citronella (*Cymbopogon nardus*, *C. winterianus*), eucalyptus (*Eucalyptus globulus*), broad and narrow-leaved eucalyptus (*E. dives* and *E. radiata*, respectively), geranium (*Pelargonium graveolens*), juniper (*Juniperus communis*), lemon (*Citrus × limon*), lemon eucalyptus (*Eucalyptus citriodora*), myrtle (*Myrtus communis*), peppermint (*Mentha × piperita*), sage (*Salvia sclarea*), thyme (*Thymus serpyllum*) verbena (*Lippia triphylla*) and wild soybean (*Glycine soja*) [128]. Differences in the species specificity of the repellence profiles of these and other EOs may explain their use in repellent mosquito products. This may have to do with differences in the local and regional profiles of mosquito species populations and explain the use of mixtures of these EOs to generate broad spec-

trum formulations for simultaneous repellency of multiple mosquito species.

While *Ae. aegypti* and other adults may in general be difficult to repel using plant-based products or synthetic repellents, a number of plant EOs have been identified which are effective against this species (► **Table 3**). The following repellency effects of EOs against adult *Ae. aegypti* have been observed: hairy basil (*Ocimum basilicum*) in stable nanoemulsions with vetiver (*Vetiveria zizanioides*) and citronella (*Cymbopogon nardus* and *C. winterianus*) EOs are a good repellent [129], bay laurel (*Laurus nobilis*) EO is an acceptable smelling, good spatial repellent [130], camphor (*Cinnamomum camphora*) EO and cassia (*C. cassia*) bark extract on human skin are repellents comparable to DEET [137], catmint (*Nepeta cataria*) EO exhibited 8 h of protection on human skin, cinnamon (*Cinnamomum zeylanicum*, *C. verum*) EOs exhibit moderate to good repellency [128], citronella (*C. winterianus*) EO exhibits knockdown repellency at 1–2% [141] and *C. winterianus* EO + vanillin exhibits 8 h of repellency [138], fennel (*Foeniculum vulgare*) fruit extract/fractions offer complete repellency [145] and EO as an aerosol or cream has comparable repellency to other EO repellents [144], geranium (*Pelargonium graveolens*) + citronella EO in a cream product exhibited good repellency in the field [46], lemon eucalyptus (*E. citriodora*) has knockdown repellency/adulticide activity [46], may chang/litsea (*Litsea cubeba*) EO exhibits contact and noncontact repellency and is a good/excellent repellent in formulations on the human forearm providing protection over 8 h [73, 128], turmeric (*Curcuma longa*) EO + 5% vanillin exhibits 8 h of repellency and in formulations with other EOs + 5% vanillin offers protection from DEET and IR3535-resistant strains of *Ae. aegypti* over 4.5 h [138, 156] and violet (*Viola odorata*) EO on human skin exhibited 8 h protection at a good level of repellency [128] (► **Table 3**).

Several EOs from *Zanthoxylum* L. spp. (*Z. piperitum* DC., *Z. armatum* DC., *Z. bungei* Planch. & Linden ex Hance) were cited in patented mosquito repellent inventions. According to recent literature, *Zanthoxylum* L. spp. EOs have mosquito repellent activity [159, 180]. Also, besides *A. argyi* H. Lév. & Vaniot (artemisia) EO, which is known to repel mosquitoes [127], the EOs of several other *Artemisia* L. spp. were used in patented formulations such as *A. annua* L. (wormwood), *A. vulgaris* L. and *A. apiacea* Hance. This is interesting given that *A. annua* EO has proven insect repellent properties [165].

The Amazon region is a source of plant-derived mosquito repellent oils. For example, *Carapa guianensis* Aubl. (andiroba) pressed fruit oils or extracts are formulated preferentially into candles during manufacturing as fumigant mosquito repellents [124], burned in kerosene lamps or used in topical repellent formulations which are commercially available in Brazil. *Copaifera* L. spp. (copaiba) balsam oils or extracts are used in mosquito repellency in Brazil especially in the Amazon region (► **Table 3**). Furthermore, a formulation of andiroba, copaiba and baby oils exhibited repellency to mosquitoes in an Amazon field study [125].

Scientific Basis for Mosquito Repellency of Glyceric Oils

▼
Neem or margarosa oil is obtained by pressing the fruit of the neem tree (*Azadirachta indica*). Neem oil is burned in 1% compositions in kerosene lamps as indoor mosquito emitters of chemical repellent-fumigant deterrents which have been evaluated and are considered to be safe [173, 174]. Also, neem oil in mixtures

Table 3 Scientific evidence for mosquito repellent and related properties of plant EOs and glyceridic oils used in patented repellent inventions.

Plant oil	Plant species	Mosquito repellent properties	Source
Essential oil			
Andiroba	<i>Carapa guianensis</i> Aubl.	patented candles commercialized in Brazil as mosquito fumigant-repellents; repellency in Amazon field test of 1 : 1 : 1 andiroba, copaiba and baby oil mixture	[124, 125]
Anise	<i>Pimpinella anisum</i> L.	good repellent against <i>Cx. pipiens</i>	[126]
Artemisia	<i>Artemisia argyi</i> H. Lévl. & Vaniot	repels mosquitoes	[127]
Basil	<i>Ocimum basilicum</i> L.	<i>Ae. aegypti</i> (PP 2 h, R% 81), <i>An. stephensi</i> (PP 3.5 h, R% 67), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; 5% hairy basil + 5% vetiver + 10% citronella EOs in nanoemulsion repel <i>Ae. aegypti</i> 4.7 h; repels <i>Anopheles</i> sp. in human bait test & <i>Cx. pipiens</i>	[128, 129]
Bay laurel	<i>Laurus nobilis</i> L.	spatial repellence against <i>Ae. aegypti</i> , acceptable smell, definite mosquito repellence	[130]
Bergamot	<i>Citrus × bergamia</i> Risso	repellence against <i>Culex</i> & <i>Aedes</i> spp. is comparable to citronella oil	[131]
Camphor	<i>Cinnamomum camphora</i> (L.) J. Presl	<i>Ae. aegypti</i> (PP 2.5 h, R% 32), <i>An. stephensi</i> (PP 8 h, R% 43), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 57)*; EO (0.1 mg · cm ⁻² on human skin): repellency and duration against <i>Ae. aegypti</i> comparable to DEET	[128, 132]
Cassia	<i>Cinnamomum cassia</i> (L.) C. Presl	bark extract (0.1 mg · cm ⁻²): repels <i>Ae. aegypti</i> comparably to DEET (on human skin), 5% EO in cream provided 50 min of protection to humans against female <i>Ae. aegypti</i>	[132]
Catnip, catmint	<i>Nepeta cataria</i> L.	<i>Ae. aegypti</i> (PP 8 h, R% 84), <i>An. stephensi</i> (PP 8 h, R% 100), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; 1–6 h protection against <i>Ae. albopictus</i> (23 & 468 µg · cm ⁻²); safety pharmacology evaluation: <i>N. cataria</i> oil is safe compared to DEET, <i>p</i> -menthane-3,8-diol, etc. May cause minor skin irritation	[128, 133, 134]
Cedar	<i>Cedrus</i> Trew (<i>Cupressus</i> L., <i>Juniperus</i> L.) spp.	<i>Ae. aegypti</i> (PP 3 h, R% 38), <i>An. stephensi</i> (PP 8 h, R% 38), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; repels <i>An. stephensi</i>	[127, 128, 135]
Chamomile	<i>Chamaemelum nobile</i> (L.) All.	<i>Ae. aegypti</i> (PP 4 h, R% 65), <i>An. stephensi</i> (PP 8 h, R% 76), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*	[128]
Cinnamon	<i>Cinnamomum zeylanicum</i> Blume	high repellency (RD ₉₅ mg · mat ⁻¹) against <i>An. stephensi</i> (49.6), <i>Ae. aegypti</i> (53.9), <i>Cx. quinquefasciatus</i> (44.2)	[136]
	<i>C. verum</i> J. Presl	<i>Ae. aegypti</i> (PP 5.5 h, R% 70), <i>An. stephensi</i> (PP 8 h, R% 100), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*	[128]
Citronella	<i>Cymbopogon nardus</i> (L.) Rendle, <i>C. winterianus</i> Jowitz ex Bor	repellency by 1 : 1 citronella & lavender EOs in 2 oz. castor oil; <i>Ae. aegypti</i> (PP 2 h, R% 76), <i>An. stephensi</i> (PP 8 h, R% 52), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; <i>C. winterianus</i> + vanillin repels <i>Ae. aegypti</i> , <i>An. dirus</i> , <i>Cx. quinquefasciatus</i> 8 h; + eucalyptus oil in floor cleaner, 6 h repellency; comparable to DEET & N,N-diethyl mandelic acid amide vs. <i>Aedes</i> spp.; 5% hairy basil + 5% vetiver + 10% citronella EOs in nanoemulsion with 4.7 h protection against <i>Ae. aegypti</i> ; <i>C. winterianus</i> LC ₅₀ & LC ₉₅ = 0.5 & 0.9% for <i>Cx. quinquefasciatus</i> , 1.0 & 2.0% for <i>Ae. aegypti</i>	[37, 128, 129, 137–141]
Clove	<i>Syzygium aromaticum</i> (L.) Merr. & L. M. Pery	<i>Cx. pipiens pallens</i> : isoeugenol & eugenol (from clove oil) > repellency than citronella. Clove bud oil + vanillin (long-term protection) > DEET; repels <i>Ae. aegypti</i> , <i>An. dims</i> , <i>Cx. quinquefasciatus</i> for 2–4 h; knockdown LC ₅₀ & LC ₉₅ = 0.5 & 0.9% for <i>Cx. quinquefasciatus</i> , 1.0 & 2.0% for <i>Ae. aegypti</i>	[46, 61, 141]
Copaiba	<i>Copaifera</i> L. spp.	1 : 1 : 1 andiroba, copaiba & baby oils repels mosquitoes in Amazon field test	[125]
Dill	<i>Anethum graveolens</i> L.	<i>Ae. aegypti</i> (PP 1.5 h, R% 78), <i>An. stephensi</i> (PP 3.5 h, R% 71), <i>Cx. quinquefasciatus</i> (PP 3 h, R% 57)*	[128]
Eucalyptus	<i>Eucalyptus globulus</i> Labill. <i>E. camaldulensis</i> Dehnh. <i>Eucalyptus</i> L'Hér. spp.	<i>Ae. aegypti</i> (PP 1 h, R% 57), <i>An. stephensi</i> (PP 5.5 h, R% 29), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)* EO repels <i>Cx. pipiens</i> adult females + citronella EO in floor cleaning product, 6 h protection; 15% oil composition ≥ 3 h protection & composition + vanillin ≥ 5 h protection to humans; EO contg 30% <i>p</i> -menthane-1,2-diol repels <i>An. darlingi</i> (97%)	[128] [8] [126, 140, 142, 143]
Eucalyptus, broad-leaved	<i>E. dives</i> Schauer	<i>Ae. aegypti</i> (PP 3.5 h, R% 19), <i>An. stephensi</i> (PP 8 h, R% 38), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*	[128]
Eucalyptus, narrow-leaved	<i>E. radiata</i> Sieber ex DC.	<i>Ae. aegypti</i> (PP 2.5 h, R% 65), <i>An. stephensi</i> (PP 8 h, R% 43), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*	[128]
Fennel	<i>Foeniculum vulgare</i> Mill.	5% in aerosol or 8% in cream repels <i>Ae. aegypti</i> comparably to citronella & geranium EOs; cream & EO repel <i>Culex</i> , <i>Anopheles</i> & <i>Aedes</i> spp. comparably to geranium & citronella EOs in field; hexane fraction (0.1 mg · cm ⁻² repels <i>Ae. aegypti</i> 99%) of fruit methanol extract contains repellent (+)-fenchone & <i>E</i> -9-octadecenoic acid	[144, 145]
Galbanum	<i>Ferula galbaniflua</i> Boiss. & Buhse	<i>Ae. aegypti</i> (PP 2.5 h, R% 70), <i>An. stephensi</i> (PP 8 h, R% 100), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*	[128]
Garlic	<i>Allium sativum</i> L.	EO and chemical components have strong repellent properties	[146]
Geranium	<i>Pelargonium graveolens</i> L'Hér.	<i>Ae. aegypti</i> (PP 2.5 h, R% 78), <i>An. stephensi</i> (PP 8 h, R% 62), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; geranium + lemongrass EOs repel <i>Aedes</i> spp.; geranium + sandalwood EOs + soybean oil in burned stick repellents; geranium (25% geraniol) EO + lemongrass extract product protect against bites of <i>Ae. atlanticus</i> and <i>Ae. Mitchellae</i> for 4 h; geranium + sandalwood EOs + soybean oil product repel <i>Culex</i> sp. & other mosquitoes for 3 h in the field; geranium + citronella EOs (1 : 1) in cream product repels <i>Ae. aegypti</i> and <i>Culex</i> , <i>Anopheles</i> & <i>Aedes</i> spp. in the field; in cold creams with repellency to <i>Cx. fatigans</i> in lab & on humans; various forms of EO definite promise as repellents	[46, 128, 137, 144, 147, 148]
Ginkgo	<i>Ginkgo biloba</i> L.	edulan (isolated from leaf extract) repels <i>Ae. albopictus</i>	[149]
Hibawood	<i>Thujopsis dolabrata</i> (Thunb. ex L. f.) Siebold & Zucc.	hinokitiol isolated from EO, highly repellent hinokitiol-containing non-woven polyester fabric & coated foot band	[150]
Jasmine	<i>Jasminum grandiflorum</i> L.	<i>Ae. aegypti</i> (PP 4.5 h, R% 14), <i>An. stephensi</i> (PP 8 h, R% 100), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*	[128]
Juniper	<i>Juniperus communis</i> L.	<i>Ae. aegypti</i> (PP 3.5 h, R% 43), <i>An. stephensi</i> (PP 8 h, R% 76), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; good repellency at 5 µg · cm ⁻² against female <i>Cx. pipiens pallens</i>	[128, 151]

continued next page

Table 3 Scientific evidence for mosquito repellent and related properties of plant EOs and glyceric oils used in patented repellent inventions. (continued)

Plant oil	Plant species	Mosquito repellent properties	Source
Essential oil			
Lavender	<i>Lavandula angustifolia</i> Mill.	<i>Ae. aegypti</i> (PP 3 h, R% 24), <i>An. stephensi</i> (PP 8 h, R% 81), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 86)*; mosquito repellent: citronella 1: lavender 1 & castor oil 2 oz.	[127, 128]
Lemon	<i>Citrus × limon</i> (L.) Osbeck	<i>Ae. aegypti</i> (PP 1.5 h, R% 68), <i>An. stephensi</i> (PP 7 h, R% 10), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; repellency to <i>An. stephensi</i> ~ DEET in animal & human tests	[128, 152]
Lemon eucalyptus	<i>Eucalyptus citriodora</i> Hook.	<i>Ae. aegypti</i> (PP 2.5 h, R% 59), <i>An. stephensi</i> (PP 8 h, R% 52), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; use (p-menthane-3,8-diol, PMD, as active ingredient) recommended by CDC (Center for Disease Control and Prevention, USA) against West Nile vector; adulticide activity against <i>Cx. quinquefasciatus</i> & <i>Ae. aegypti</i>	[8, 46, 128]
Lemongrass	<i>Cymbopogon citratus</i> (DC.) Stapf	<i>Ae. aegypti</i> (PP 3 h, R% 70), <i>An. stephensi</i> (PP 8 h, R% 100), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; field test of 25% geraniol oil & lemongrass ext. with 4 h repellency against <i>Ae. atlanticus</i> and <i>Ae. mitchellae</i> ; mixture containing lemongrass EO + p-menthane diol (PMD) repels <i>An. darlingi</i> & other spp. 95–98% for 5–6 h in field (repellency > 15–20% DEET)	[128, 147, 153]
Lemon tea tree	<i>Leptospermum petersonii</i> F. M. Bailey	limited repellency (< DEET) of <i>Ae. aegypti</i> , <i>Cx. quinquefasciatus</i> , <i>Cx. annulirostris</i>	[154]
Leptospermum	<i>L. liversidgei</i> R. T. Baker & H. G. Sm.	repellent: blocks ability of mosquitoes to perceive CO ₂ emitted by humans	[155]
Lime	<i>Citrus × aurantifolia</i> L.	+ mustard oil carrier as effective mosquito repellent	[68]
Marjoram	<i>Origanum majorana</i>	5 µg · cm ⁻² good repellent against female <i>Cx. pipiens pallens</i>	[151]
May chang/ Litsea	<i>Litsea cubeba</i> (Lour.) Pers.	<i>Ae. aegypti</i> (PP 8 h, R% 73), <i>An. stephensi</i> (PP 8 h, R% 100), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; high contact and noncontact repellency to female <i>Ae. aegypti</i> <i>in vitro</i> in cages; formulation to fix on skin with high repellency in human volunteers for up to 8 h & 100% repellency to <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i> , <i>Ae. aegypti</i> ; greater repellence of night-biting mosquitoes <i>An. dirus</i> , <i>Cx. quinquefasciatus</i> and <i>Ae. albopictus</i> than <i>Ae. aegypti</i>	[73, 128, 132, 156]
Melaleuca	<i>Melaleuca alternifolia</i> Cheel	limited repellence of <i>Ae. aegypti</i> , <i>Cx. quinquefasciatus</i> , <i>Cx. annulirostris</i> in lab	[75]
Mint, mentha	<i>Mentha haplocalyx</i> Briq.	d-8-acetoxycarvotanacetone isolated mosquito repellent	[157]
Myrtle	<i>Myrtus communis</i> L.	<i>Ae. aegypti</i> (PP 2.5 h, R% 57), <i>An. stephensi</i> (PP 6.5 h, R% 43), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 86)*	[128]
Palmarosa	<i>Cymbopogon martini</i> (Roxb.) Will. Watson	high geraniol content mosquito repellent	[158]
Patchouli	<i>Pogostemon cablin</i> (Blanco) Benth.	partially repels <i>Ae. aegypti</i> , <i>Cx. quinquefasciatus</i> , <i>An. dirus</i>	[159]
Pepper, black	<i>Piper nigrum</i> L.	<i>Ae. aegypti</i> (PP 1.5 h, R% 65), <i>An. stephensi</i> (PP 3 h, R% 62), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; repels <i>An. dirus</i> , <i>Cx. quinquefasciatus</i> , <i>Ae. albopictus</i> (≥ 4.5 h)	[128, 156]
Peppermint	<i>Mentha × piperita</i> L.	<i>Ae. aegypti</i> (PP 2 h, R% 59), <i>An. stephensi</i> (PP 6.5 h, R% 57), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; on human skin strongly repels <i>An. annularis</i> (100%), <i>An. culicifacies</i> (92%), <i>Cx. quinquefasciatus</i> (85%); comparable to myl oil (di-butyl & di-methyl phthalates); repels adult female <i>Cx. pipiens</i>	[126, 128, 160]
Pine	<i>Pinus sylvestris</i> L.	good mosquito repellency	[85]
Rose	<i>Rosa</i> L. sp.	moderate mosquito repellency	[127]
Rosemary	<i>Rosmarinus officinalis</i> L.	<i>Ae. aegypti</i> (PP 5.5 h, R% 43), <i>An. stephensi</i> (PP 8 h, R% 100), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; repels <i>An. stephensi</i> , <i>Ae. aegypti</i> & <i>Cx. quinquefasciatus</i> ; low repellency against <i>Ae. aegypti</i> in lab, acceptable to humans as final fragrance in repellent formulation; repels <i>Ae. aegypti</i> (avg 90 min)	[88, 128, 130, 136]
Sage	<i>Salvia sclarea</i> L.	<i>Ae. aegypti</i> (PP 2 h, R% 46), <i>An. stephensi</i> (PP 5 h, R% 19), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*	[128]
Sandalwood	<i>Santalum album</i> L. <i>Santalum</i> L. spp.	<i>Ae. aegypti</i> (PP 2.5 h, R% 59), <i>An. stephensi</i> (PP 8 h, R% 100), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)* repels <i>Cx. pipiens pallens</i> ; EO containing cold creams repel <i>Cx. fatigans</i> in the lab & on human skin; area repellency of <i>Culex</i> sp. by commercial sticks (contg 0.5% EO) (3 h protection); botanical repellent (w/soybean and geranium oil) 100% effective in field (3-h test period, comparable to DEET)	[128] [93, 148, 151]
Sour (bitter) orange	<i>Citrus × aurantium</i> L.	limited protection (repellency) against mosquitoes	[137]
Spearmint	<i>Mentha spicata</i> L.	repellent; piperitone oxide from <i>M. spicata</i> (var. <i>viridis</i>) EO repels <i>An. stephensi</i>	[161]
Tagetes	<i>Tagetes minuta</i> L.	<i>Ae. aegypti</i> (PP 1 h, R% 84), <i>An. stephensi</i> (PP 8 h, R% 100), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; repels <i>Ae. aegypti</i> (avg 90 min)	[88, 128]
Thyme	<i>Thymus serpyllum</i> L. <i>Thymus vulgaris</i> L.	<i>Ae. aegypti</i> (PP 2.5 h, R% 57), <i>An. stephensi</i> (PP 7.5 h, R% 33), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)* repels lab-reared adult female <i>Ae. albopictus</i> for 2 h; 0.01% totally repels <i>Cx. quinquefasciatus</i>	[128] [134, 162]
Turmeric	<i>Curcuma longa</i> L.	+ 5% vanillin repels <i>Ae. aegypti</i> , <i>An. dirus</i> and <i>Cx. quinquefasciatus</i> in cage & large room for 8 h; turmeric, citronella & hairy basil EOs + vanillin provide substitute for DEET; repels DEET & IR3535-resistant <i>Ae. aegypti</i> for 4.5 h	[138, 156]
Verbena	<i>Lippia triphylla</i> (L'Hér.) Kuntze	<i>Ae. aegypti</i> (PP 2.5 h, R% 70), <i>An. stephensi</i> (PP 5.5 h, R% 38), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*	[128]
Vetiver	<i>Vetiveria zizanioides</i> (L.) Nash	stable nanoemulsion of 5% hairy basil, 5% vetiver & 10% citronella EO: repels <i>Ae. aegypti</i> for 4.7 h	[41]
Violet	<i>Viola odorata</i> L.	<i>Ae. aegypti</i> (PP 6 h, R% 68), <i>An. stephensi</i> (PP 8 h, R% 100), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 86)*	[128]
Wild verbena (Lippie)	<i>Lippia sidoides</i> , <i>L. javanica</i> , <i>Lippia</i> spp.	sources of perillaldehyde (repels <i>An. gambiae</i>) & perillic acid (repels <i>An. arabiensis</i> & <i>Ae. aegypti</i>)	[163, 164]
Wintergreen	<i>Gaultheria procumbens</i> L.	strongly repels <i>Culex</i> and <i>Aedes</i> spp. in different tests	[131]
Wormwood	<i>Artemisia annua</i> L.	mosquito repellent properties	[165]

continued next page

Table 3 Scientific evidence for mosquito repellent and related properties of plant EOs and glyceridic oils used in patented repellent inventions. (continued)

Plant oil	Plant species	Mosquito repellent properties	Source
Essential oil			
Zanthoxylum	<i>Z. piperitum</i> DC., <i>Z. armatum</i> DC., <i>Z. bungei</i> Planch. & Linden ex Hance	repel mosquitoes	[159]
Glyceridic oils			
Castor	<i>Ricinus communis</i> L.	best carrier for pyrethrum extracts (long-lasting)	[137]
Mustard	<i>Brassica</i> L. spp.	longer protection (up to 5 h with <i>Zanthoxylum limonella</i> or lime oils) than coconut (<i>Cocos nucifera</i>) oil against <i>Ae. albopictus</i>	[166]
Neem/ Margosa	<i>Azadirachta indica</i> A. Juss.	knockdown repellency against <i>Ae. aegypti</i> , <i>Ae. albopictus</i> , <i>An. quadrimaculatus</i> Say; repels female <i>An. stephensi</i> (ED ₅₀ 0.191–0.156 mg · cm ⁻²) in lab; low repellency against <i>Ae. albopictus</i> & <i>Cx. nigripalpus</i> ; 2% in coconut (<i>Cocos nucifera</i>) oil on exposed body parts of human volunteers provided complete protection for 12 h from all <i>Anopheles</i> spp.; protection from <i>Anopheles</i> spp. (96–100%), <i>Aedes</i> (85%), <i>Culex</i> sp. (61–94%); significant protection by neem cream against adult <i>Ae. aegypti</i> ; 1% in kerosene lamps in preclinical & clinical safety evaluation is safe to humans; 1% in kerosene burned in lamps effective in 2 field tests. Repellence: <i>Anopheles</i> > <i>Culex</i> ; 1–4% in coconut oil on exposed body parts of humans: 81–91% protection for 12 h	[167–175]
Olive	<i>Olea europaea</i> L.	<i>Ae. aegypti</i> (PP 3.5 h, R% 68), <i>An. stephensi</i> (PP 8 h, R% 71), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 71)*; 1:1 w/pyrethrum repels mosquitoes for 4 h	[128, 137]
Soybean	<i>Glycine max</i> (L.) Merr.	<i>Ae. aegypti</i> (PP 3 h, R% 54), <i>An. stephensi</i> (PP 8 h, R% 76), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*; oil-based product provided 1.5 h (low) repellency, 24% DEET provided 5 h protection; good repellency in product w/sandalwood + geranium in burned sticks vs. DEET; commercial product containing 2% oil repels <i>Ae. albopictus</i> , <i>Cx. nigripalpus</i> , <i>Ae. triseriatus</i> for 5, 8.5 & ≥ 7.3 h, respectively	[128, 169, 176, 177]
Soybean, wild (carotin)	<i>Glycine soja</i> Siebold & Zucc.	<i>Ae. aegypti</i> (PP 3 h, R% 16), <i>An. stephensi</i> (PP 8 h, R% 10), <i>Cx. quinquefasciatus</i> (PP 8 h, R% 100)*	[128]

* PP = protection period (h), R% = percent repellency; 0.1 mL of a 20% EO solution (in 20% Genapol, 10% PEG, 20% ethanol, 50% water) applied to 30 cm² of human forearm; for 20% DEET solution (in same as above) these values were: *Ae. aegypti* (PP 6 h, R% 46), *An. stephensi* (PP 8 h, R% 100), *Cx. quinquefasciatus* (PP 8 h, R% 100) [24]

with coconut (*Cocos nucifera* L.) oil provides good protection for very long periods against *Anopheles* and *Aedes* spp. [170, 171, 175]. Castor (*Ricinus communis* L.), mustard (*Brassica* spp.), olive (*Olea europaea*) and other glyceridic oils have important roles in several patented repellent compositions containing pyrethrum extracts and EOs where they act as carriers and can extend the duration of repellent effects for several hours perhaps by slowing the release or evaporation of EOs from surfaces.

There are mixed scientific reports on the effective mosquito repellency of several glyceridic oils. Thus, for soybean (*Glycine max*) oil low repellency was observed as compared to 24% DEET formulations [176] and good repellency was observed for the smoke generated from burned sticks which contained soybean oil. Especially interesting is a report from a United States Department of Agriculture laboratory where 4 well-known synthetic mosquito repellents based on 10% KBR3023 [1-piperidinecarboxylic acid 2-(2-hydroxyethyl)-1-methylpropyl ester], 7.5% IR3535 [3-(*N*-butyl-*N*-acetyl)-aminopropionic acid ethyl ester], 15% and 7% DEET and 8 natural product-based repellents based on 2% soybean oil, 10% citronella (*Cymbopogon* spp.) EO, neem oil (*Azadirachta indica*) and others were tested in the lab against *Ae. albopictus* Skuse, *Cx. nigripalpus* Theobald and *Ae. triseriatus* Say [169]. The 2% soybean oil formulation exhibited mosquito repellency comparable to both 10% KBR3023 and 15% DEET based products each of which provided estimated mean protection time (eMPT) responses averaged over all three mosquito species of ≥ 7.2 h [169]. This study is evidence for the potential of soybean oil as a stand-alone repellent and as a component oil of repellent formulations.

Mosquito Repellent Chemical Components of EOs and Added Isolated Compounds

Approximately 20% of all EO-containing patents mentioned a non-EO plant derivative and 40% mentioned a specific EO repellent component or added isolated natural chemical ingredient. Chrysanthamic acid and its derivatives which are components of pyrethrum (*Chrysanthemum* L. spp. flower extracts) were cited in 12% of all patented formulations involving EOs. Also, patents cited many “natural” chemical constituents which were either added to formulations or were important chemical components of EOs comprising the formulations. The most widely cited chemicals of plant origin were (% of patents): camphor (8.3), geraniol (7.6), linalool (7.6), menthol (7.6), geranial (7.0), citronellal (6.9), limonene (4.9), citronellol (4.9), borneol (4.2), 1,8-cineole (4.2), *p*-menthane-3,8-diol (4.2), camphene (3.5) and nepetalactone and derivatives (3.5). Many of these and other chemical components of EOs are potent mosquito repellents (Table 4) [131, 137, 143, 145, 151, 163, 169, 181–193] and are discussed below.

Vanillin in EO containing mosquito repellent inventions

Vanillin was cited as an additive in 4% of EO-containing patents. According to several scientific publications, it increases the duration and magnitude of the repellent effect of synthetic repellents such as DEET as well as plant EOs. For example, 5% vanillin in formulations of EOs of turmeric (*Curcuma longa* L.), citronella (*C. winterianus*) or hairy basil (*Ocimum americanum* L.) provided up to 8 h of protection against *Ae. aegypti*, *An. dirus* and *Cx. quinquefasciatus* under cage conditions. Interestingly, 5% vanillin in formulations with DEET increased the protection time as compared to DEET alone against these three mosquito species (≥ 8 h protec-

Table 4 Mosquito repellency or detergency effects of individual major chemical components of EOs used in patented inventions.

EO component	No. EOs [†]	Repellency	Mosquito	Source
α -Bisabolol	2	^d 84% spatial repellency for 180 min, avoidance freq. (contact repellency) = DEET	female <i>Ae. aegypti</i>	[181]
Borneol	3	^b 1.4 mg (-)-borneol · cm ⁻² = 70% repellency ^{b,c} RD ₅₀ = 1.7 × 10 ⁻³ mg · cm ⁻²	female <i>Ae. aegypti</i> female <i>An. gambiae</i>	[182] [163]
Bornyl acetate	2	^b 1.4 mg · cm ⁻² = 15% repellency	female <i>Ae. aegypti</i>	[182]
Camphene	2	^{b,c} RD ₅₀ = 2.2 × 10 ⁻³ mg · cm ⁻²	female <i>An. gambiae</i>	[163]
Camphor	6	^{b,c} RD ₅₀ = 1.4 × 10 ⁻³ mg · cm ⁻² ^b 1.4 mg · cm ⁻² (\pm)-camphor, (+)-camphor, (-)-camphor = 36, 35 & 47% repellencies, resp.	female <i>An. gambiae</i> female <i>Ae. aegypti</i>	[163] [182]
3-Carene	4	^{b,c} RC ₅₀ = 8.6 × 10 ⁻⁴ mg · cm ⁻² 1 h, 1.92 μ g · cm ⁻² = 68 ^f , 65 ^g % repellency	female <i>An. gambiae</i> <i>Ae. aegypti</i> , <i>Ae. albopictus</i>	[183] [163]
Carvacrol	4	^{b,c} RC ₅₀ = 2.4 × 10 ⁻⁴ mg · cm ⁻²	female <i>An. gambiae</i>	[183]
Carvone	2	^{b,c} RD ₅₀ = 1.3 × 10 ⁻³ mg · cm ⁻² <i>d</i> -carvone & <i>l</i> -carvone: feeding deterrents/repellents, good spatial repellents/ inhibitors, contact repellents/deterrents	female <i>An. gambiae</i> -	[163] [184]
Cinnamaldehyde	2	definite mosquito repellent potential <i>E</i> -cinnamaldehyde (0.051 mg · cm ⁻²) & DEET (0.025 mg · cm ⁻²) provided 87 & 95% protection, resp., after 30 min	- female <i>Aedes aegypti</i>	[137] [184]
1,8-Cineole	15	^b cineole 1.4 mg · cm ⁻² = 21% repellency ^b 3% in vaseline or olive oil = 73 or 120 min protection ^{b,c} RD ₅₀ = 1.2 × 10 ⁻³ mg · cm ⁻²	female <i>Ae. aegypti</i> 1 : 1 male : female <i>Cx. pipiens molestus</i> female <i>An. gambiae</i>	[182] [185] [163]
Citronellal	4	^{b,c} RD ₅₀ = 2.2 × 10 ⁻⁴ mg · cm ⁻² feeding deterrent/repellent, spatial repellent/inhibitor, contact repellent/deterrent	female <i>An. gambiae</i> -	[163] [186]
Citronellol	6	repellent component feeding deterrent/repellent, spatial repellent/inhibitor	- -	[145] [186]
<i>p</i> -Cymene	3	^{b,c} RD ₅₀ = 1.0 × 10 ⁻⁵ mg · cm ⁻²	female <i>An. gambiae</i>	[163]
Eugenol	5	high repellency ^{b,c} RD ₅₀ = 1.3 × 10 ⁻³ mg · cm ⁻²	female <i>Cx. pipiens pallens</i> female <i>An. gambiae</i>	[151] [163]
Fenchone	2	^{b,c} RD ₅₀ = 1.9 × 10 ⁻³ mg · cm ⁻² 0.4 mg (+)-fenchone · cm ⁻² = 51% repellency after 1 h ^{b,f}	female <i>An. gambiae</i> female <i>Ae. aegypti</i>	[163] [145]
Geranial	7	at 0.2 mg · cm ⁻² , blood-sucking by mosquitoes reduced by 10, 15, and 18% after 1, 2 and 3 h, respectively	<i>Cx. pipiens pallens</i> , <i>Cx. pipiens</i> <i>quinquefasciatus</i>	[187]
Geraniol	11	^{b,c} RD ₅₀ = 1.1 × 10 ⁻⁴ mg · cm ⁻² spatial repellent: 0.25 μ g · cm ⁻² ca. 100% (reversible) inhibition of host-seeking after 48 h exposure	female <i>An. gambiae</i> female <i>Ae. albopictus</i>	[163] [188]
Geranyl acetate	6	effective feeding deterrent/repellent, effective spatial repellent/inhibitor, contact repellent/deterrent	Mosquitoes	[186]
Limonene	19	^b 3% in vaseline/olive oil = 61/78 min protection 1 h, 1.92 μ g · cm ⁻² = 70 ^f , 70 ^g % repellency ^{b,c} RD ₅₀ = 1.8 × 10 ⁻³ mg · cm ⁻²	1 : 1 male : female <i>Cx. pipiens molestus</i> <i>Ae. aegypti</i> , <i>Ae. albopictus</i> female <i>An. gambiae</i>	[185] [189] [163]
Linalool	14*	^b 1.4 mg (\pm)-linalool · cm ⁻² = 67% repellency ^b 3% in vaseline/olive oil = 52/65 min protection ^{b,c} RD ₅₀ = 1.5 × 10 ⁻³ mg · cm ⁻²	female <i>Ae. aegypti</i> 1 : 1 male : female <i>Cx. pipiens molestus</i> female <i>An. gambiae</i>	[182] [185] [163]
<i>p</i> -Menthane-3,8-diol	1	26% formula more repellent than 4 synthetic & 7 natural product repellents 15% formula with lemongrass EO (<i>Cymbopogon citratus</i>) = 99% protection for 5 h in the field; ^h 16% formula with lemongrass EO = 95% protection for 6 h in the field	<i>Ae. albopictus</i> , <i>Cx. nigripalpus</i> , <i>Ochlerotatus triseriatus</i> <i>An. darling</i> , <i>Ae. ochlerotatus</i> <i>taeniorhynchus</i>	[169] [143]
Menthone	2	^b 3% in olive oil = 35 min protection	1 : 1 male : female <i>Cx. pipiens molestus</i>	[185]
Methyl salicylate	2	most repellent (screening)	-	[131]
Myrcene	6	^{b,c} RC ₅₀ = 8.4 × 10 ⁻⁴ mg · cm ⁻² ^b 1.4 mg · cm ⁻² = 20% repellency	female <i>An. gambiae</i> <i>Ae. aegypti</i>	[183] [182]
Nepetalactone	1	mosquito repellent ingredient in formulations	-	[190]
<i>E</i> -Nerolidol	1	^d 67% spatial repellency for 180 min, avoidance freq. (contact repellency) < DEET	female <i>Ae. aegypti</i>	[181]
Oleic acid	8**	0.4 mg · cm ⁻² = 52% repellency after 1 h ^{b,f}	female <i>Ae. aegypti</i>	[145]
Perillaldehyde	1	^{b,c} RD ₅₀ = 3.2 × 10 ⁻⁴ mg · cm ⁻²	female <i>An. gambiae</i>	[163]
α -Pinene	15	<i>d</i> - α -pinene, 2 h protection ^b 3% in vaseline/olive oil = 27/56 min protection	Mosquito 1 : 1 male : female <i>Cx. pipiens molestus</i>	[191], [192] [185]
β -Pinene	8	^b 3% in vaseline/olive oil = 22/39 min protection ^{b,c} RD ₅₀ = 1.6 × 10 ⁻³ mg · cm ⁻²	1 : 1 male : female <i>Cx. pipiens molestus</i> female <i>An. gambiae</i>	[185] [163]
α -Santalol	1	^d 87% spatial repellency for 180 min, avoidance freq. (contact repellency) = DEET	female <i>Ae. aegypti</i>	[181]
Spathulenol	2	^e proportion not biting = 0.73, 0.75, respectively	<i>Ae. aegypti</i> , <i>An. stephensi</i>	[193]

continued next page

Table 4 Mosquito repellency or deterrence effects of individual major chemical components of EOs used in patented inventions. (continued)

EO component	No. EOs [†]	Repellency	Mosquito	Source
α-Terpinene	1	1 h, 1.92 μg · cm ⁻² = 68 [†] , 55 [‡] % repellency	<i>Ae. aegypti</i> , <i>Ae. albopictus</i>	[189]
		^b 1.4 mg · cm ⁻² = 15 % repellency	female <i>Ae. aegypti</i>	[182]
		^{b,c} RD ₅₀ = 2.4 × 10 ⁻³ mg · cm ⁻²	female <i>An. gambiae</i>	[163]
γ-Terpinene	6	1 h, 1.92 μg · cm ⁻² = 60 [†] , 72 [‡] % repellency	<i>Ae. aegypti</i> , <i>Ae. albopictus</i>	[189]
		^b 3% in vaseline/olive oil = 35/48 min protection	1 : 1 male : female <i>Cx. pipiens molestus</i>	[185]
		^{b,c} RD ₅₀ = 2.7 × 10 ⁻³ mg · cm ⁻²	female <i>An. gambiae</i>	[163]
Terpinen-4-ol	5	1 h, 1.92 μg · cm ⁻² = 92 [†] , 85 [‡] % repellency	<i>Ae. aegypti</i> , <i>Ae. albopictus</i>	[189]
		^b 1.4 mg · cm ⁻² = 83 % repellency	<i>Ae. aegypti</i>	[182]
		^{b,c} RD ₅₀ = 1.5 × 10 ⁻³ mg · cm ⁻²	female <i>An. gambiae</i>	[163]
		spatial repellent/inhibitor	–	[186]
α-Terpineol	7	^b 3% in vaseline/olive oil = 78/99 min protection	1 : 1 male : female <i>Cx. pipiens molestus</i>	[185]
		feeding deterrent/repellent, spatial repellents/inhibitor, contact repellent/deterrent		[186]
		^{b,c} RD ₅₀ = 1.3 × 10 ⁻³ mg · cm ⁻²	female <i>An. gambiae</i>	[163]
Terpinolene	2	1 h, 1.92 μg · cm ⁻² = 55 [†] , 70 [‡] % repellency	<i>Ae. aegypti</i> , <i>Ae. albopictus</i>	[189]
		^{b,c} RD ₅₀ = 2.6 × 10 ⁻³ mg · cm ⁻²	female <i>An. gambiae</i>	[163]
(α + β)-Thujone	2	^b 1.4 mg · cm ⁻² = 20 % repellency	<i>Ae. aegypti</i>	[182]
Thymol	1	^{b,c} RC ₅₀ = 1.9 × 10 ⁻³ mg · cm ⁻²	female <i>An. gambiae</i>	[183]
		^b 3% in vaseline/olive oil = 52/65 min protection	1 : 1 male : female <i>Cx. pipiens molestus</i>	[186]
cis-Verbenol	1	^{b,c} RD ₅₀ = 7.5 × 10 ⁻⁵ mg · cm ⁻²	female <i>An. gambiae</i>	[163]
Verbenone	1	^{b,c} RD ₅₀ = 1.6 × 10 ⁻³ mg · cm ⁻²	female <i>An. gambiae</i>	[163]

– = Information not found. [†] Number of oils in **Table 1** which contain this as their major component. *In evening primrose (*Oenothera biennis*) glyceridic oil. ** Mosquito repellent component of glyceridic oils. [†] DEET = 50 % repellency. [‡] DEET = 60 % repellency. ^a Used as additive in one patent. ^b Human-bait assay. ^c Positive control DEET exhibited RC₅₀ = 3.3 × 10⁻⁴ mg · cm⁻². ^d Applied concentration 78.6 μg · cm⁻². Repellency is statistically different from negative control. DEET and turmerone had 83 and 89 % repellencies, respectively. Contact repellency is defined as 100 % of individuals off treated surface. Avoidance frequency of DEET and turmerone = 0.8. ^e Substance was applied at 25 nmol · cm⁻² cloth. Positive control was (1S,2'S)-2-methylpiperidinyl-3-cyclohexen-1-carboxamide (SS-220) at 25 nmol · cm⁻² cloth (provided 0.80 and 0.78 non-biting mosquitoes, respectively, for *Ae. aegypti* and *An. stephensi*). ^f 0.2 mg · cm⁻² DEET (positive control) provided 97 % repellency after 1 h. ^g Positive control 15 % DEET (in EtOH) = 92 % protection (average over trial period).

^h Positive control 20 % DEET (in EtOH) = 64 % protection (average over trial period)

tion). In another example, 15% eucalyptus (*Eucalyptus* spp.) EO and 5% vanillin provided 5 h of repellency against *Ae. albopictus* [18] and in other work involving human volunteers, clove (*Syzygium aromaticum*) bud EO and vanillin mixtures provided long acting repellency against adult female *Cx. pipiens pallens* [151]. Thus, combinations of vanillin with plant EOs can lead to formulations which are substitutes for DEET [138, 142, 151].

In patents, vanillin is cited as an ingredient in mosquito repellent formulations with: *Zanthoxylum armatum* DC. and/or *Z. piperitum* DC. EOs [194], clove bud and leaf (*Syzygium aromaticum*), juniper berry (*Juniperus communis*) and/or marjoram (*Origanum majorana*) EOs [195], lemongrass (*Cymbopogon citratus*) EO/p-menthane-3,8-diol formulations [196], combinations of citronella, clove, geranium (*Pelargonium graveolens*), lavender (*Lavandula angustifolia*), patchouli [*Pogostemon cablin* (Blanco) Benth.] and peppermint (*Mentha × piperita*) EOs [197] among others.

Synthetic additives in EO-containing inventions

About 25% of all EO-containing patented repellents had at least one synthetic repellent component. Thus, DEET (*N,N*-diethyl *m*-toluamide), the gold-standard synthetic mosquito repellent [11], was used in 10% and dimethyl (dialkyl) phthalates were cited in 5% of EO-containing patented inventions. Also, one or more synthetic pyrethroids (allethrin, cyhalothrin, deltamethrin, dimefluthrin, esbiothrin, metofluthrin, permethrin, tetramethrin and vaporthrin) were used in 10% of EO-containing mosquito repellents. It must be stressed that pyrethroids can be strong mosquitoicides (toxicity), spatial mosquito repellents and mosquito irritants as was shown, for example, for permethrin in field studies [6] and α-cypermethrin, deltamethrin, permethrin in a very re-

cent publication [198]. Only about 3% of patents made use of the synthetic synergist piperonyl butoxide.

Chemical Composition of EOs Used in Patented Inventions

Information on the major chemical components of plant EOs used in mosquito repellent inventions is presented in **Table 1**. Among these are EOs which are concentrated sources of proven mosquito repellent monoterpenes and phenylpropanoids such as the EOs of angelica (*Angelica archangelica*), artemisia (*Artemisia argyi*), basil (*Ocimum basilicum*), bergamot (*Citrus × bergamia*), camphor (*Cinnamomum camphora*), cassia (*Cinnamomum cassia*), catnip (*Nepeta cataria*), chrysanthemum (*Chrysanthemum indicum*), cinnamon (*Cinnamomum zeylanicum*), citronella (*Cymbopogon nardus*, *C. winterianus*), coriander (*Coriandrum sativum*), cypress (*Cupressus sempervirens*), dill (*Anethum graveolens*), eucalyptus (*Eucalyptus* spp.), geranium (*Pelargonium graveolens*), grapefruit (*Citrus reticulata*), ho leaf (*Cinnamomum camphora*), hyssop (*Hyssopus officinalis*), juniper (*Juniperus communis*), lavender (*Lavandula angustifolia*), lemon (*Citrus × limon*), lemon eucalyptus (*Eucalyptus citriodora*), lemongrass (*Cymbopogon citratus*), lemon tea tree (*Leptospermum petersonii*), lime (*Citrus × aurantifolia*), marjoram (*Origanum majorana*), may chang/litsea (*Litsea cubeba*), melaleuca/tea tree (*Melaleuca alternifolia*), mint/mentha (*Mentha*), orange (*Citrus × sinensis*), palmarosa (*Cymbopogon martini*), curl leaf parsley (*Petroselinum crispum*), pepper (*Piper nigrum*), pine (*Pinus sylvestris*), rose (*Rosa × damascena*, *R. × centifolia*), rosemary (*Rosmarinus officinalis*), salvia/sage (*Salvia* spp.), sour (bitter) orange (*Citrus × aurantium*),

spearmint (*Mentha spicata*), thyme (*Thymus vulgaris*), verbena (*Lippia triphylla*) and wintergreen (*Gaultheria procumbens*). In these EOs, proven mosquito repellent volatile components camphor, 1,8-cineole, citronellol, eugenol, geranial, geraniol, limonene, linalool, myrcene, α and β -pinenes, γ -terpinene, terpinen-4-ol and α -terpineol are well represented among the major components (Table 4).

Chemical composition of glyceridic oils used in patented EO-containing inventions

Glyceridic plant oils are important components of EO-containing mosquito repellent formulations not only for their carrier properties, but because they contain trace or larger amounts of free, long-chain mono- and polyunsaturated fatty acids and derivatives of these. It has been known for some time that insect antennae are highly sensitive to gas-phase long-chain fatty acids. For example, Hwang et al. [199] showed that gravid *Cx. quinquefasciatus* Say in an olfactometer in contact with room temperature vapors of C₁₄-C₂₄ mono-, di- and polyunsaturated fatty acids were significantly repelled. These authors also demonstrated that fatty acids having at least one Z-configuration olefin bond (analogous saturated and E-configuration fatty acids were less repellent or inactive) were especially effective repellents. One of the most repellent fatty acids was oleic acid (9Z-olefin) which is a common free fatty acid in vegetable oils whose 18 carbon chain was considered to be optimal within the group of fatty acids studied [199]. Independently, female *Ae. aegypti* were found to be moderately repelled by oleic acid as compared to DEET in a human-bait assay [145]. Thus, free, long-chain, unsaturated fatty acids which are present in glyceridic oils are sufficiently volatile to be mosquito repellent substances in formulations containing plant EOs. This is especially relevant given claims in several patents of the mosquito repellent effects of olive (*Olea europaea* L.), sesame (*Sesamum indicum* L.), mustard (*Brassica* L. spp.), soybean [*Glycine max* (L.) Merr.] and other oils (for examples, see [200] or [201]).

Mosquito repellent major chemical components of EOs

Among the more widely distributed mosquito repellent major chemical components of EOs used in patented inventions are the cyclic and noncyclic monoterpenes limonene, α -pinene, 1,8-cineole and linalool (Table 4). These are major components of, respectively, 28, 22 [including limonene-containing perilla (*Perilla frutescens*) oil], 22 and 21% [including linalool-containing evening primrose (*Oenothera biennis*) oil] of the EOs most used in patented repellent inventions (Table 1).

EOs in patented mosquito repellent inventions are composed of a number of chemical components which individually exhibit important mosquito repellency, deterrence and inhibitory effects (Table 1 and Table 4). For example, α -bisabolol and α -santalol provided 3 h of high spatial repellency and contact repellency equivalent to DEET against female *Ae. aegypti* while E-nerolidol exhibited good spatial repellency for 3 h and less contact repellency than DEET [181]. Also, the following substances singly exhibited good repellency against the mosquito species indicated: carvacrol (RC₅₀ = 2.4 × 10⁻⁴ mg·cm⁻², female *An. gambiae*) [183], cinnamaldehyde (0.051 mg·cm⁻² against female *Aedes aegypti*, repellency comparable to DEET) [184], citronellal (RD₅₀ = 2.2 × 10⁻⁴ mg·cm⁻²), *p*-cymene (RD₅₀ = 1.0 × 10⁻⁵ mg·cm⁻²) and *cis*-verbenol (RD₅₀ = 7.5 × 10⁻⁵ mg·cm⁻²) against female *An. gambiae* [163], geranial (0.2 mg·cm⁻², reduces blood-sucking by *Cx. pipiens pallens* and *Cx. pipiens quinquefasciatus* over 3 h). The fol-

lowing EO components provided reasonable repellency at ca. 2 µg·cm⁻² against adult *Ae. aegypti* and *Ae. albopictus*: 3-carene, limonene, α -terpinene, γ -terpinene, terpinen-4-ol and terpinolene [189].

Several interesting isolated repellent substances are described in patented EO-containing products (Table 3). For example, hinokitiol-containing polyester non-woven fabric and hinokitiol-coated polyester foot band are said to provide 100% repellency against mosquitoes and hinokitiol from hibawood (*Thujaopsis dolabrata*) EO (and other sources) is an important natural repellent [150]. Another example is the isolation of edulan from ginkgo (*Ginkgo biloba*) EO and the repellent activity of this compound and derivatives against *Ae. albopictus* are described in a patent in which it is also claimed that these compounds are safe to humans [149]. The mosquito repellent natural substance *d*-8-acetoxycarvotanacetone was isolated from *Mentha haplocalyx* Briq. [157]. Also, several *Lippia* spp. EOs were cited in patents [e.g., *L. sidoides* and *L. javanica* (wild verbena or lippie oil)] and are important sources of the mosquito repellent compounds perillaldehyde and perillaldehyde [163, 164] (Table 3). The latter compound is a component of perilla (*Perilla frutescens*) oil and exhibited good repellency (RD₅₀ = 3.2 × 10⁻⁴ mg·cm⁻²) against female *An. gambiae* [163] (Table 4). In another patented invention, garlic (*Allium sativum*) EO and isolated component chemicals of garlic EO were tested for repellency against mosquitoes. More than a dozen allyl sulfide, allyl disulfide and allyl polysulfide components of garlic EO applied to human skin were found to deter *Ae. aegypti*, *An. gambiae* and *Cx. quinquefasciatus* landing and blood feeding [202].

Geraniol

Geraniol is a major component of 16% of the more used EOs in patented inventions (Table 1) and its mosquito repellency has been studied. As an isolated chemical, it exhibited good repellency on human skin in the lab [163] and reversible spatial repellency/protection from bites [188]. A 25% formulation of this substance was the basis for a commercial product which exhibited effective repellency in the lab [93, 167, 169] but was less effective as compared to other products at repelling mosquitoes according to an early field study [143]. In a recent field study, a 25% geraniol and lemongrass (*Cymbopogon citratus*) composition offered superior protection against the bites of *Ae. atlanticus* and *Ae. mitchellae* (4 h) than repellent formulations based on 12% EOs (1 h) and EOs, benzophenone-3, octinoxate and octisalate mixture (1.5 h) [147].

p-Menthane-3,8-diol

The mosquito repellency of the component of *Eucalyptus* spp. EOs, *p*-menthane-3,8-diol (PMD), is firmly established [8, 143, 178]. As a repellent additive in formulations, 15–16% PMD/lemongrass formulations provided 5–6 h of excellent repellency (95–99%) against *Anopheles* and *Aedes* spp. [153]. Indeed, in a U.S. Dept. of Agriculture study, 26% PMD-containing product was more repellent in the lab to *Ae. albopictus*, *Cx. nigripalpus* and *Ae. triseriatus* than synthetic products based on 10% KBR3023, 7.5% IR3535, 15% and 7% DEET, and 7 natural product-based repellents based on 2% soybean oil, 10% citronella EO, neem oil, 25% geraniol, etc. [169].

Mosquito repellent minor components of EOs

Some minor or lower concentration components of EOs are worth mentioning because they exhibit significant mosquito repellency. For example, citral (geranial + neral) and fenchyl alcohol are minor components of citronella (*Cymbopogon nardus*, *C. winterianus*) and rose (*Rosa damascena*) EOs, respectively, and exhibit moderate repellency to female *An. gambiae* ($RD_{50} = 1.3 \times 10^{-3}$ and $1.4 \times 10^{-3} \text{ mg} \cdot \text{cm}^{-2}$, respectively) [163]. Also, linalool oxide, a minor component of lemon eucalyptus (*E. citriodora*) EO was found to be repellent ($RC_{50} = 6.5 \times 10^{-4} \text{ mg} \cdot \text{cm}^{-2}$) to female *An. gambiae* [183] and Tripathi et al. [161] found that the peppermint (*Mentha × piperita*) EO component piperitone oxide was highly repellent to adult *An. stephensi*. In other works, farnesol [a component of chamomile, jasmine and rose (*Rosa damascena*) EOs], elemol (a component of catnip, citronella and hyssop EOs) and hedycaryol (a component of black pepper EO) exhibited 69, 89 and 95% spatial repellency over 180 min to adult female *Ae. aegypti* and avoidance frequency (contact repellency) greater than or equal to that of DEET [181]. Cinnamyl alcohol, a minor constituent of cassia (*Cinnamomum cassia*) EO, exhibits less repellency to female *Ae. aegypti* than the major component, cinnamaldehyde, but still this repellency is significant (at $0.051 \text{ mg} \cdot \text{cm}^{-2}$, 86% protection after 30 min) and worth mentioning here [184]. Also, a minor component of EO of catmint (*Nepeta* L. spp.) EOs [28], dihydronepetalactone, exhibits important repellency against arthropods, including mosquitoes [203]. Citronellyl acetate is a minor component of eucalyptus (*Eucalyptus* spp.) [165], lemon eucalyptus (*E. citriodora*) [66], rose (*Rosa damascena*) [87] and tarragon (*Artemisia dracuncululus*) [99] EOs and is a feeding deterrent/repellent and mosquito spatial repellent/inhibitor. Also, the minor component of salvia/sage EO [92], hydroxycymene, is both a feeding deterrent/repellent and mosquito contact repellent/deterrent [186]. Lastly, isoeugenol (a component of clove EO) was found to be highly repellent to female *Cx. pipiens pallens* [151].

Synthetic EOs: Additive Effects/Synergism versus Suppression/Dilution

In some cases, patented formulations in effect improve upon the natural repellency of an essential oil by mimicking or synthesizing an oil which ideally contains components which together contribute to the repellent effect and eliminates those which counteract (attract mosquitoes), make no contribution or suppress the repellency of other components. Work done by Odalo et al. [183] nicely illustrates this process. An initial observation was that major components of 6 EOs when tested singly were less repellent than the natural EOs in which these components are found. Synthetic EOs were prepared by mixing pure major components in the same proportion in which they occur in the natural EOs. Repellencies of synthetic EOs ranged from comparable to up to three times the activity of the corresponding natural EOs against *An. gambiae* in the human-bait test based on RC_{75} values [183]. The activity of synthetic EOs substantiates the additive and/or synergistic nature of the interaction of blended EO components (and also suppressive or diluting/repellency diminishing effects of nonactive components). The same principles of addition/synergism and suppression/dilution are operational in the process of formulating EOs and isolated components into patented repellent inventions.

Enantiomeric Composition of EOs

Scientific (and patent) literature on the composition of EOs is often based on non-chiral column gas chromatography-mass spectrometry (GC-MS) and compound retention time/index analysis. Thus, information is not systematically available on the enantiomeric purity of the EO components in **Table 1**. Enantio-specificity in mosquito repellency is an important issue. For example, Gu et al. [189] found that commercial *R*-(-)-terpinen-4-ol exhibited >90% repellency against *Ae. aegypti* and *Ae. albopictus* adults at a concentration of $1.92 \mu\text{g} \cdot \text{cm}^{-2}$ in a test lasting 1 h. *S*-(+)-terpinen-4-ol of high enantiomeric purity is found in lavender (*Lavandula angustifolia*) EOs [204]. No mosquito repellency data were found in the literature for *S*-(+)-terpinen-4-ol. In principle, enantiomers may not be equally repellent to mosquitoes and their organoleptic and other properties (toxic, allergic, etc.) are not equally desirable for use as repellents. More research is needed on the comparative mosquito repellency of enantiomerically pure EO components such as *S*-(+)-terpinen-4-ol and *R*-(-)-terpinen-4-ol against different mosquito species.

Allergenicity of EO Chemical Components

Limonene, benzyl alcohol, linalool, citronellol, geraniol, citral, anisyl alcohol, cinnamaldehyde, eugenol, isoeugenol, coumarin, farnesol, benzyl benzoate, benzyl salicylate among other volatile components are known allergens. As described above, these components are major or minor components of EOs used in repellent inventions (**Table 1** and references cited therein). These and other volatile allergens are easily detected by GC-MS and are controlled in the European Union and elsewhere.

Conclusion

Patents on plant EO and volatile component-containing mosquito repellent inventions make use of scientifically substantiated claims to repellency to mosquitoes based on lab and field experiences. Our experience is that mosquito repellents and other insect control products, must be evaluated in the environments and settings in which they are to be used. Repellents must be developed based on plant oils and isolated chemical components by targeting *Aedes*, *Anopheles*, *Culex* and other mosquito species which are found locally and regionally. Repellency approaches should be multipronged and make use of sprays, fumigants, paints and varnishes, incense, candles, etc. in domestic settings, fumigation and spraying in outdoor settings and topical repellents, clothes made of repellent fabrics, repellent wristbands among other available products for individual protection.

Acknowledgements

The authors are grateful to Joanne Bero and Joëlle Quetin-Leclercq for the invitation for this review. Our thanks also go to the anonymous referee for many invaluable corrections and suggestions. The authors are thankful for grants and/or scholarships received from Brazilian agencies MCT/INPA, CNPq 561559/2008-2 and 555.669/2009-2 (Brazilian Malaria Network), CNPq Bionorte Network grant 2009, FAPERN.

References

- 1 WHO Media Centre. Malaria. World Health Organization (WHO) Media Centre. Available at <http://www.who.int/mediacentre/factsheets/fs094/en/>. Accessed September 26, 2010.
- 2 WHO Media Centre. Dengue. World Health Organization (WHO) Media Centre. Available at <http://www.who.int/mediacentre/factsheets/fs117/en/>. Accessed September 26, 2010.
- 3 Burfield T, Reekie SL. Mosquitoes, malaria and essential oils. *Int J Aromather* 2005; 15: 30–41.
- 4 WHO Media Centre. Filariasis. World Health Organization (WHO) Media Centre. Available at <http://www.who.int/topics/filariasis/en/>. Accessed September 26, 2010.
- 5 WHO Media Centre. Yellow Fever. World Health Organization (WHO) Media Centre. Available at <http://www.who.int/mediacentre/factsheets/fs100/en/>. Accessed September 26, 2010.
- 6 Brown M, Hebert AA. Insect repellents: an overview. *J Am Acad Dermatol* 1997; 36: 243–249.
- 7 Nerio LS, Olivero-Verbel J, Stashenko E. Repellent activity of essential oils: a review. *Bioresour Technol* 2010; 101: 372–378.
- 8 Batish DR, Singh HP, Kohli RK, Kaur S. Eucalyptus essential oil as a natural pesticide. *For Ecol Manage* 2008; 256: 2166–2174.
- 9 EPA. US Environmental Protection Agency. Notice to manufacturers, formulators, producers and registrants of pesticide products on minimum risk pesticides exempted under FIFRA Section 25(b) Pesticide Registration (PR). Notice 2000–6, 2000. Labeling Unit, Registration Division (7505C). Washington; Office of Pesticide Programs, EPA. 2000. Available from: http://www.epa.gov/pesticides/biopesticides/ingredients/factsheets/factsheet_plant-oils.htm. Accessed September 26, 2010.
- 10 Bakkali F, Averbeck S, Averbeck D, Zhiri A, Idaomar M. Cytotoxicity and gene induction by some essential oils in the yeast *Saccharomyces cerevisiae*. *Mutat Res* 2005; 585: 1–13.
- 11 SciFinder Scholar. Chemical Abstracts Service – A division of the American Chemical Society. Available at <http://www.cas.org/products/sf-cad/index.html>. Accessed September 28, 2010.
- 12 Tropicos.org. Missouri botanical garden. Available at <http://www.tropicos.org>. Accessed November 30, 2010.
- 13 Rout PK, Rao YR, Jena KS, Sahoo D, Mishra BC. Extraction and composition of the essential oil of ambrette (*Abelmoschus moschutos*) seeds. *J Essent Oil Res* 2004; 16: 35–37.
- 14 Rout PK, Barik KC, Jena KS, Sahoo D, Rao YR. A novel process for the extraction of fragrance components from ambrette (*Hibiscus abelmoschus* L.) seeds. *Org Proc Res Dev* 2002; 6: 401–404.
- 15 Nivinskiene O, Butkiene R, Mockute D. Changes in the chemical composition of essential oil of *Angelica archangelica* L. roots during storage. *Chemija (Vilnius)* 2003; 14: 52–56.
- 16 Wu M, Sun X, Dai Y, Guo F, Huang L, Liang Y. Determination of constituents of essential oil from *Angelica sinensis* by gas chromatography-mass spectrometry. *J Central South Univ Technol* 2005; 12: 430–436.
- 17 Arslan N, Gürbüz B, Sarihan EO, Bayrak A, Gümüşçü A. Variation in essential oil content and composition in Turkish anise (*Pimpinella anisum* L.) populations. *Turkish J Agric For* 2004; 28: 173–177.
- 18 Wenqiang G, Shufen L, Ruixiang Y, Yanfeng H. Comparison of composition and antifungal activity of *Artemisia argyi* Lévl. et Vant inflorescence essential oil extracted by hydrodistillation and supercritical carbon dioxide. *Nat Prod Res* 2006; 20: 992–998.
- 19 Zheng X, Deng C, Song G, Hu Y. Comparison of essential oil composition of *Artemisia argyi* leaves at different collection times by headspace solid-phase microextraction and gas chromatography-mass spectrometry. *Cromatographia* 2004; 59: 729–732.
- 20 Inouye S, Uchida K, Abe S. Volatile composition and vapour activity against *Trichophyton mentagrophytes* of 36 aromatic herbs cultivated in Chichibu district in Japan. *Int J Aromather* 2006; 16: 159–168.
- 21 Viña A, Murillo E. Essential oil composition from twelve varieties of basil (*Ocimum* spp.) grown in Colombia. *J Braz Chem Soc* 2003; 14: 744–749.
- 22 Chalchat J-C, Özcan MM. Comparative essential oil composition of flowers, leaves and stems of basil (*Ocimum basilicum* L.) used as herb. *Food Chem* 2008; 110: 501–503.
- 23 Sangun MK, Aydin E, Timur M, Karadeniz H, Caliskan M, Ozkan A. Comparison of chemical composition of the essential oil of *Laurus nobilis* L. leaves and fruits from different regions of Hatay, Turkey. *J Environment Biol* 2007; 28: 731–733.
- 24 Cosimi S, Rossi E, Cioni PL, Canale A. Bioactivity and qualitative analysis of some essential oils from Mediterranean plants against stored-product pests: evaluation of repellency against *Sitophilus zeamais* Motschulsky, *Cryptolestes ferrugineus* (Stephens) and *Tenebrio molitor* (L.). *J Stored Prod Res* 2009; 45: 125–132.
- 25 Wang R, Wang R, Yang B. Extraction of essential oils from five cinnamon leaves and identification of their volatile compound compositions. *Innovative Food Sci Emerg Technol* 2009; 10: 289–292.
- 26 Du H, Li H. Antioxidant effect of cassia essential oil on deep-fried beef during the frying process. *Meat Sci* 2008; 78: 461–468.
- 27 Gilani AH, Shaha AJ, Zubair A, Khalid S, Kiani J, Ahmed A, Rasheed M, Ahmad VU. Chemical composition and mechanisms underlying the spasmodic and bronchodilatory properties of the essential oil of *Nepeta cataria* L. *J Ethnopharmacol* 2009; 121: 405–411.
- 28 Heuskun S, Godin B, Leroy P, Capella Q, Wathelet J-P, Verheggen F, Haubruge E, Lognay G. Fast gas chromatography characterisation of purified semiochemicals from essential oils of *Matricaria chamomilla* L. (Asteraceae) and *Nepeta cataria* L. (Lamiaceae). *J Chromatogr A* 2009; 1216: 2768–2775.
- 29 Burfield T. Cedarwood oils. Part 1. *Aromather Times* 2002; 1: 14–15.
- 30 Lawrence BM. Progress in essential oils. *Perfumer Flavorist* 2005; 30: 56–67.
- 31 Wu L-Y, Gao H-Z, Wang X-L, Ye J-H, Lu J-L, Liang Y-R. Analysis of chemical composition of *Chrysanthemum indicum* flowers by GC/MS and HPLC. *J Med Plant Res* 2010; 4: 421–426.
- 32 Zhu S, Yang Y, Yu H, Ying Y, Zou G. Chemical composition and antimicrobial activity of the essential oils of *Chrysanthemum indicum*. *J Ethnopharmacol* 2005; 96: 151–158.
- 33 Fichi G, Flamini G, Zaralli LJ, Perrucci S. Efficacy of an essential oil of *Cinnamomum zeylanicum* against *Psoroptes cuniculi*. *Phytomedicine* 2007; 14: 227–231.
- 34 Singh G, Maurya S, DeLampasona MP, Catalan CAN. A comparison of chemical, antioxidant and antimicrobial studies of cinnamon leaf and bark volatile oils, oleoresins and their constituents. *Food Chem Toxicol* 2007; 45: 1650–1661.
- 35 Nakahara K, Alzoreky NS, Yoshihashi T, Nguyen HTT, Trakoontivakorn G. Chemical composition and antifungal activity of essential oil from *Cymbopogon nardus* (citronella grass). *Jpn Intl Res Center Agric Sci* 2003; 37: 249–252.
- 36 Sethi ML, Taneja SC, Dhar KL, Atal CK. Chemical investigations on the essential oil of *Burmese citronella*. *Indian Perfumer* 1979; 23: 167–171.
- 37 Guan W, Li S, Yan R, Tang S, Quan C. Comparison of essential oils of clove buds extracted with supercritical carbon dioxide and other three traditional extraction methods. *Food Chem* 2007; 101: 1558–1564.
- 38 Msaada K, Taarit MB, Hosni K, Hammami M, Marzouk B. Regional and maturational effects on essential oils yields and composition of coriander (*Coriandrum sativum* L.) fruits. *Sci Hortic* 2009; 122: 116–124.
- 39 Zheljzkova VD, Pickett KM, Caldwell CD, Pincok JA, Roberts JC, Mapplebeck L. Cultivar and sowing date effects on seed yield and oil composition of coriander in Atlantic Canada. *Indian Crops Prod* 2008; 28: 88–94.
- 40 Tapondjou AL, Adler C, Fontem DA, Bouda H, Reichmuth C. Bioactivities of cymol and essential oils of *Cupressus sempervirens* and *Eucalyptus saligna* against *Sitophilus zeamais* Motschulsky and *Tribolium confusum* du Val. *J Stored Prod Res* 2005; 41: 91–102.
- 41 Sacchetti OG, Maietti S, Muzzoli M, Scaglianti M, Manfredini S, Radice M, Bruni R. Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. *Food Chem* 2005; 91: 621–632.
- 42 Bailer J, Aichinger T, Hackl G, de Hueber K, Dachler M. Essential oil content and composition in commercially available dill cultivars in comparison to caraway. *Indian Crops Prod* 2001; 14: 229–239.
- 43 Callan NW, Johnson DL, Westcott MP, Welty LE. Herb and oil composition of dill (*Anethum graveolens* L.): Effects of crop maturity and plant density. *Indian Crops Prod* 2007; 25: 282–287.
- 44 Kimbaris AC, Siatis NG, Daferera DJ, Tarantilis PA, Pappas CS, Polissiou MG. Comparison of distillation and ultrasound-assisted extraction methods for the isolation of sensitive aroma compounds from garlic (*Allium sativum*). *Ultrason Sonochem* 2006; 13: 54–60.
- 45 Shaath NA, Flores FB, Ositian M, Aal MA-E. The essential oil of *Allium sativum* L., Liliaceae (garlic). In: *Charalambous G*, editor. *Food flavors: generation, analysis and process influence*. Amsterdam: Elsevier Science; 1995: 2025–2037.

- 46 Seo S-M, Kim J, Lee S-G, Shin C-H, Shin S-C, Park I-K. Fumigant antitermitic activity of plant essential oils and components from ajowan (*Trachyspermum ammi*), allspice (*Pimenta dioica*), caraway (*Carum carvi*), dill (*Anethum graveolens*), geranium (*Pelargonium graveolens*), and litsea (*Litsea cubeba*) oils against Japanese termite (*Reticulitermes speratus* Kolbe). *J Agric Food Chem* 2009; 57: 6596–6602
- 47 Singh G, Kapoor IPS, Singh P, de Heluani CS, de Lampasona MP, Catalan CAN. Chemistry, antioxidant and antimicrobial investigations on essential oil and oleoresins of *Zingiber officinale*. *Food Chem Toxicol* 2008; 46: 3295–3302
- 48 Sakamura F. Changes in volatile constituents of *Zingiber officinale* rhizomes during storage and cultivation. *Phytochemistry* 1987; 26: 2207–2212
- 49 Chutia M, Deka Bhuyan P, Pathak MG, Sarma TC, Boruah P. Antifungal activity and chemical composition of *Citrus reticulata* Blanco essential oil against phytopathogens from North East India. *LWT – Food Sci Technol* 2009; 42: 777–778
- 50 Bates RB, Slagel RC. Terpenoids. VI. β -Bulnesene, α -guaiene, β -patchoulene, and guaioxide in essential oils. *Chem Ind* 1962; 1715–1716
- 51 Yatagai M, Sat T, Takahashi T. Terpenes of leaf oils from Cupressaceae. *Biochem Syst Ecol* 1985; 13: 377–385
- 52 Burfield T. Cedarwood oils – part 2. *Aromather Times* 2003; 1: 14–16
- 53 Nozoe T, Yasue A, Yamane K. On the acidic constituents of the essential oil of *Thujopsis dolabrata*. Occurrence of α -thujaplicin. *Proc Jpn Acad* 1951; 27: 15–17
- 54 Zhang G, Chen C, Chen Z, Chen R, Lin X. Analysis of principle component and chemistry type of essential oil from *Cinnamomum camphora* leaf in Fujian province. *Zhiwu Ziyuan Yu Huanjing Xuebao* 2008; 17: 24–27
- 55 Nguyen XD, Trinh DC, Nguyen BT, Pham VK, Leclercq PA. Constituents of the leaf oil of *Cinnamomum camphora* Nees et Eberm from Hue. *Tap Chi Hoa Hoc* 1994; 32: 64–66
- 56 Kazazi H, Rezaei K, Ghotb-Sharif SJ, Emam-Djomeh Z, Yamini Y. Supercritical fluid extraction of flavors and fragrances from *Hyssopus officinalis* L. cultivated in Iran. *Food Chem* 2007; 105: 805–811
- 57 Langa E, Cacho J, Palavra AMF, Burillo J, Mainar AM, Urieta JS. The evolution of hyssop oil composition in the supercritical extraction curve. *J Supercrit Fluids* 2009; 49: 37–44
- 58 Chalchat J-C, Adamovic D, Gorunovic MS. Composition of oils of three cultivated forms of *Hyssopus officinalis* endemic in Yugoslavia: f. *albus* Alef., f. *cyaneus* Alef. and f. *ruber* Mill. *J Essent Oil Res* 2001; 13: 419–421
- 59 Rath CC, Devi S, Dash SK, Mishra RK. Antibacterial potential assessment of jasmine essential oil against *E. coli*. *Indian J Pharm Sci* 2008; 70: 238–241
- 60 Sandeep, Paarakh PM. *Jasminum grandiflorum* Linn (Chameli): ethnobotany, phytochemistry and pharmacology – a review. *Pharmacol Online* 2009; 2: 586–595
- 61 Angioni A, Barra A, Russo MT, Coroneo V, Dessi S, Cabras P. Chemical composition of the essential oils of *Juniperus* from ripe and unripe berries and leaves and their antimicrobial activity. *J Agric Food Chem* 2003; 51: 3073–3078
- 62 Al-Reza SM, Rahman A, Kang SC. Chemical composition and inhibitory effect of essential oil and organic extracts of *Cestrum nocturnum* L. on food-borne pathogens. *Int J Food Sci Technol* 2009; 44: 1176–1182
- 63 Salehi FARP, Heydari R, Ebrahimi SN, Haddad PR. Hydrodistillation-headspace solvent microextraction, a new method for analysis of the essential oil components of *Lavandula angustifolia* Mill. *J Chromatogr A* 2005; 1098: 14–18
- 64 Daferera DJ, Ziogas BN, Polissiou MG. GC-MS analysis of essential oils from some Greek aromatic plants and their fungitoxicity on *Penicillium digitatum*. *J Agric Food Chem* 2000; 48: 2576–2581
- 65 Flamini G, Tebano M, Cioni PL. Volatiles emission patterns of different plant organs and pollen of *Citrus limon*. *Anal Chim Acta* 2007; 589: 120–124
- 66 El-Zalabani SM, Koheil MM, Meselhy KM, El-Gizawy HA, Sleem AA. Effect of seasonal variation on composition and bioactivities of the essential oil of *Eucalyptus citriodora* Hook. grown in Egypt. *Egyptian J Biomed Sci* 2007; 24: 260–276
- 67 Brophy JJ, Goldsack RJ, Punruckvong A, Bean AR, Forster PI, Lepschi BJ, Doran JC, Rozefelds AC. Leaf essential oils of the genus *Leptospermum* (Myrtaceae) in eastern Australia. Part 7. *Leptospermum petersonii*, *L. liversidgei* and allies. *Flav Fragr J* 2000; 15: 342–351
- 68 Patil JR, Jayaprakasha GK, Murthy KNC, Tichy SE, Chetti MB, Patil BS. Apoptosis-mediated proliferation inhibition of human colon cancer cells by volatile principles of *Citrus aurantifolia*. *Food Chem* 2009; 114: 1351–1358
- 69 Douglas MH, van Klink JW, Smallfield BM, Perry NB, Anderson RE, Johnston, Weavers RT. Essential oils from New Zealand manuka: triketone and other chemotypes of *Leptospermum scoparium*. *Phytochemistry* 2004; 65: 1255–1264
- 70 Perry NB, Brennan NJ, Van Klink JW, Harris W, Douglas MH, McGimpsey JA, Smallfield BM, Anderson RE. Essential oils from New Zealand manuka and kanuka: chemotaxonomy of *Leptospermum*. *Phytochemistry* 1997; 44: 1485–1494
- 71 Vági E, Simándi B, Suhajda A, Héthelyi E. Essential oil composition and antimicrobial activity of *Origanum majorana* L. extracts obtained with ethyl alcohol and supercritical carbon dioxide. *Food Res Int* 2005; 38: 51–57
- 72 Vera RR, Chane-Ming J. Chemical composition of the essential oil of marjoram (*Origanum majorana* L.) from Reunion Island. *Food Chem* 1999; 66: 143–145
- 73 Lalko J, Api AM. Investigation of the dermal sensitization potential of various essential oils in the local lymph node assay. *Food Chem Toxicol* 2006; 44: 739–746
- 74 Lawrence BM. Progress in essential oils. *Perfumer Flavorist* 2006; 31: 54–57
- 75 Brophy JJ, Davies NW, Southwell IA, Stiff IA, Williams LR. Gas chromatographic quality control for oil of melaleuca terpinen-4-ol type (Australian tea tree). *J Agric Food Chem* 1989; 37: 1330–1335
- 76 Marengo E, Baiocchi C, Gennaro MC, Bertolo PL, Lanteri S, Garrone W. Classification of essential mint oils of different geographic origin by applying pattern recognition methods to gas chromatographic data. *Chromometrics Intell Lab Syst* 1991; 11: 75–88
- 77 Tao N-G, Liu Y-J, Zhang M-L. Chemical composition and antimicrobial activities of essential oil from the peel of bingtang sweet orange (*Citrus sinensis* Osbeck). *Int J Food Sci Technol* 2009; 44: 1281–1285
- 78 Raina VK, Srivastava SK, Aggarwal KK, Syamasundar KV, Khanuja SPS. Essential oil composition of *Cymbopogon martinii* from different places in India. *Flav Fragr* 2003; 18: 312–315
- 79 Petropoulos SA, Daferera D, Polissiou MG, Passam HC. Effect of nitrogen-application rate on the biomass, concentration, and composition of essential oils in the leaves and roots of three types of parsley. *J Plant Nutr Soil Sci* 2009; 172: 210–215
- 80 Nikiforov A, Jirovetz L, Buchbauer G, Raverdino V. GC-FTIR and GC-MS in odour analysis of essential oils. *Microchim Acta* 2005; 95: 193–198
- 81 Orav A, Stulova I, Kailas T, Müürisepp M. Effect of storage on the essential oil composition of *Piper nigrum* L. fruits of different ripening states. *J Agric Food Chem* 2004; 52: 2582–2586
- 82 Kapoor IPS, Singh B, Singh G, De Heluani CS, De Lampasona MP, Catalan CAN. Chemistry and *in vitro* antioxidant activity of volatile oil and oleoresins of black pepper (*Piper nigrum*). *J Agric Food Chem* 2009; 57: 5358–5364
- 83 Kowalski R, Wawrzykowski J. Essential oils analysis in dried materials and granulates obtained from *Thymus vulgaris* L., *Salvia officinalis* L., *Mentha piperita* L. and *Chamomilla recutita* L. Essential oil analysis in dried materials and granulates. *Flav Fragr* 2009; 24: 31–35
- 84 Aziz EE, Craker LE. Essential oil constituents of peppermint, pennyroyal, and apple mint grown in a desert agrosystem. *J Herbs Spices Med Plants* 2009; 15: 361–367
- 85 Judzentiene A, Kupcinskiene E. Chemical composition on essential oils from needles of *Pinus sylvestris* L. grown in northern Lithuania. *J Essent Oil Res* 2008; 20: 26–29
- 86 Ustun O, Sezik E, Kurkcuoglu M, Baser KHC. Study of the essential oil composition of *Pinus sylvestris* from Turkey. *Chem Nat Comp* 2006; 42: 26–31
- 87 Almasirad A, Amanzadeh Y, Taheri A, Iranshahi M. Composition of a historical rose oil sample (*Rosa damascena* Mill., Rosaceae). *J Essent Oil Res* 2007; 19: 110–112
- 88 Gillij YG, Gleiser RM, Zygodlo JA. Mosquito repellent activity of essential oils of aromatic plants growing in Argentina. *Bioresour Technol* 2008; 99: 2507–2515
- 89 Geneva MP, Stancheva IV, Boychinova MM, Mincheva NH, Yonova PA. Effects of foliar fertilization and arbuscular mycorrhizal colonization on *Salvia officinalis* L. growth, antioxidant capacity, and essential oil composition. *J Sci Food Agric* 2010; 90: 696–702

- 90 Kamatou GPP, Makunga NP, Ramogola WPN, Viljoen AM. South African *Salvia* species: a review of biological activities and phytochemistry. *J Ethnopharmacol* 2008; 119: 664–672
- 91 Loizzo MR, Menichini F, Tundis R, Bonesi M, Nadjafi F, Saab AM, Frega NG, Menichini F. Comparative chemical composition and antiproliferative activity of aerial parts of *Salvia lerifolia* Benth. and *Salvia acetabulosa* L. essential oils against human tumor cell *in vitro* models. *J Med Food* 2010; 13: 62–69
- 92 Cardile V, Russo A, Formisano C, Rigano D, Senatore F, Arnold NA, Piozzi F. Essential oils of *Salvia bracteata* and *Salvia rubifolia* from Lebanon: chemical composition, antimicrobial activity and inhibitory effect on human melanoma cells. *J Ethnopharmacol* 2009; 126: 265–272
- 93 Jones CG, Plummer JA, Barbour EL. Non-destructive sampling of Indian sandalwood (*Santalum album* L.) for oil content and composition. *J Essent Oil Res* 2007; 19: 157–164
- 94 Lawrence BM. Progress in essential oils. *Perfumer Flavorist* 2005; 30: 56–67
- 95 Kiran U, Patra DD. Influence of natural essential oils and their by-products as nitrification retarders in regulating nitrogen utilization for Japanese mint in sandy loam soils of subtropical central India. *Agric Ecosyst Environ* 2003; 94: 237–245
- 96 Wang Z, Wang L, Li T, Zhou X, Ding L, Yu Y, Yu A, Zhang H. Rapid analysis of the essential oils from dried *Illicium verum* Hook. f. and *Zingiber officinale* Rosc. by improved solvent-free microwave extraction with three types of microwave-absorption medium. *Anal Bioanal Chem* 2006; 386: 1863–1868
- 97 Dzamic A, Sokovic M, Ristic MS, Grijic-Jovanovic S, Vukojevic J, Marin PD. Chemical composition and antifungal activity of *Illicium verum* and *Eugenia caryophyllata* essential oils. *Chem Nat Comp* 2009; 45: 259–261
- 98 Lawrence BM. Progress in essential oils. *Perfumer Flavorist* 2009; 34: 54–58
- 99 Khodakov GV, Kotikov IV, Pankovetskii VN. Component composition of essential oil from *Artemisia abrotanum* and *A. dracunculus*. *Chem Nat Comp* 2009; 45: 905–908
- 100 Argyropoulou C, Daferera D, Tarantilis PA, Fasseas C, Polissiou M. Chemical composition of the essential oil from leaves of *Lippia citriodora* H.B.K. (Verbenaceae) at two developmental stages. *Biochem Syst Ecol* 2007; 35: 831–837
- 101 Champagnat P, Figueredo G, Chalchat J-C, Carnat A-P. A study on the composition of commercial *Vetiveria zizanioides* oils from different geographical origins. *J Essent Oil Res* 2006; 18: 416–422
- 102 Cu JQ, Perineau F, Gaset A. Volatile components of violet leaves. *Phytochemistry* 1992; 31: 571–573
- 103 Tisserand R. Essential oil safety. *Int J Aromather* 1996; 7: 28–32
- 104 Martín C, Moure A, Martín G, Carrillo E, Domínguez H, Parajó JC. Fractional characterisation of jatropha, neem, moringa, trisperma, castor and candlenut seeds as potential feedstocks for biodiesel production in Cuba. *Bioresour Technol* 2010; 34: 533–538
- 105 Bicchì C, Cordero C, Liberto E, Sgorbini B, Rubiolo P. Headspace sampling of the volatile fraction of vegetable matrices. *J Chromatogr A* 2008; 1184: 220–233
- 106 Christie WW. The analysis of evening primrose oil. *Indian Crops Prod* 1999; 10: 73–83
- 107 Jeong JB, Ju SY, Park JH, Lee JR, Yun KW, Kwon ST, Lim J-H, Chung GY, Jeong HJ. Antioxidant activity in essential oils of *Cnidium officinale* makino and *Ligusticum chuanxiong* hort and their inhibitory effects on DNA damage and apoptosis induced by ultraviolet B in mammalian cell. *Cancer Epidemiol* 2009; 33: 41–46
- 108 Nabloussi A, Márquez-Lema M, Fernández-Martínez A, Velasco L. Novel seed oil types of Ethiopian mustard with high levels of polyunsaturated fatty acids. *Indian Crops Prod* 2008; 27: 359–363
- 109 Sidhu OP, Kumar V, Behl HM. Variability in triterpenoids (nimbin and salanin) composition of neem among different provenances of India. *Indian Crops Prod* 2004; 19: 69–75
- 110 Patumi M, d'Andria R, Marsilio V, Fontanazza G, Morelli G, Lanza B. Olive and olive oil quality after intensive monocone olive growing (*Olea europaea* L., cv. Kalamata) in different irrigation regimes. *Food Chem* 2002; 77: 27–34
- 111 Nitta M, Kobayashi H, Ohnishi-Kameyama M, Nagamine T, Yoshida M. Essential oil variation of cultivated and wild perilla analyzed by GC/MS. *Biochem Syst Ecol* 2006; 34: 25–37
- 112 Longvah T, Deosthale YG, Kumar PU. Nutritional and short term toxicological evaluation of *Perilla* seed oil. *Food Chem* 2000; 70: 13–16
- 113 Koezuka Y, Honda G, Tabata M. Genetic control of the chemical composition of volatile oils in *Perilla frutescens*. *Phytochemistry* 1986; 25: 859–863
- 114 Lee D-S, Lee E-S, Kim H-J, Kim S-O, Kim K. Reversed phase liquid chromatographic determination of triacylglycerol composition in sesame oils and the chemometric detection of adulteration. *Anal Chim Acta* 2001; 429: 321–330
- 115 Ribeiro APB, Grimaldi R, Gioielli LA, Gonçalves LAG. Zero trans fats from soybean oil and fully hydrogenated soybean oil: physico-chemical properties and food applications. *Food Res Int* 2009; 42: 401–410
- 116 Orban GJ, Staton JA. Insect repellent compositions based on essential oils. *World Patent* 2004010783; 2004
- 117 He J, Yang X, Wang Y, Li W, Zhao J, Feng J, Zhang X. Plant-derived mosquito repellent synergist. *Chinese Patent* 101558769; 2009
- 118 Samson GB. Organic floor wash with disinfectant and mosquito repellent properties. *Indian Patent* 2007CH00296; 2007
- 119 Wang Z, Wan T, Zhang J, Liu H, Zhang X, Feng J, Zhang X. Plant-derived synergistic agent for mosquito incense and its preparation process. *Chinese Patent* 101558770; 2009
- 120 Inoue S, Hario S, Umibe H. Synergistic insect-repelling microcapsules containing toluamides and isobornyl compounds, for textiles. *Japanese Patent* 03148203; 1991
- 121 Peng X. Mosquito-repellent solution containing natural pyrethrin. *Chinese Patent* 1739353; 2006
- 122 Bit KC, Rao SR, Sen K. Synergistic insect repellent compositions containing dimethyl phthalate, allyl sulfate, and oils. *Indian Patent* 2003DE00243; 2007
- 123 Singh KSP, Kumar TA, Kumar AK, Kumar SM, Kumar SA, Raj BJ, Anju Y, Dinesh K, Sudeep T, Anirban P, Umrao BD, Mansoor A, Prakash DO. Slow-vaporizing anti-mosquito synergistic insecticidal herbal formulation comprising essential oil. *Indian Patent* 2008DE00607; 2010
- 124 Gilbert B, Teixeira DF, Pereira JFG, Silva Jr IV, Svaiter A, Oliveira Filho AM, Santos CE. Mosquito-repellent fumigant. *Brazilian Patent* 9800437; 2000
- 125 Ribas J, Carreno AM. Evaluation of the use of repellent against mosquito bite by military personnel in the Amazon Basin. *An Bras Dermatol* 2010; 85: 33–38
- 126 Erler F, Ulug I, Yalcinkaya B. Repellent activity of five essential oils against *Cx. pipiens*. *Fitoterapia* 2006; 77: 491–494
- 127 Lin X. Repellent effects of plant essential oils on laboratory animals. *Zhonghua Weisheng Shachong Yaoxue* 2009; 15: 430–431
- 128 Amer A, Mehlhorn H. Repellency effect of forty-one essential oils against *Aedes*, *Anopheles*, and *Culex* mosquitoes. *Parasitol Res* 2006; 99: 478–490
- 129 Onanong N, Usawadee S, Napaporn U, Satit P, Apinan S, Uracha R. *In vitro* characterization and mosquito (*Aedes aegypti*) repellent activity of essential-oils-loaded nanoemulsions. *AAPS Pharm Sci Technol* 2009; 10: 1234–1242
- 130 Drapeau J, Froehler C, Touraud D, Kroeckel U, Geier M, Rose A, Kunz W. Repellent studies with *Aedes aegypti* mosquitoes and human olfactory tests on 19 essential oils from Corsica, France. *Flav Fragr J* 2009; 24: 160–169
- 131 Rudolfs W. Effect of chemicals upon the behavior of mosquitoes. *New Jersey Agric Exp Station Bull* 1930; 2: 496–520
- 132 Yang Y-C, Lee E-H, Lee H-S, Lee D-K, Ahn Y-J. Repellency of aromatic medicinal plant extracts and a steam distillate to *Aedes aegypti*. *J Am Mosq Control Assoc* 2004; 20: 146–149
- 133 Zhu J, Zeng X, Yanma LT, Qian K, Han Y, Xue S, Tucker B, Schultz G, Coats J, Rowley W, Zhang A. Adult repellency and larvicidal activity of five plant essential oils against mosquitoes. *J Am Mosq Control Assoc* 2006; 22: 515–522
- 134 Zhu JJ, Zeng X-P, Berkebile D, Du H-J, Tong Y, Qian K. Efficacy and safety of catnip (*Nepeta cataria*) as a novel filth fly repellent. *Med Vet Entomol* 2009; 23: 209–216
- 135 Curtis CF, Lines JD, Callaghan A, Hill N, Karimzad MA. The relative efficiency of repellents against mosquito vectors of disease. *Med Vet Entomol* 1987; 1: 109–119
- 136 Prajapati V, Tripathi AK, Aggarwal KK, Khanuja SPS. Insecticidal, repellent and oviposition-deterrent activity of selected essential oils against *Anopheles stephensi*, *Aedes aegypti* and *Cx. quinquefasciatus*. *Bioresour Technol* 2005; 96: 1749–1757
- 137 Gibson A. Mosquito-suppression work in Canada in 1937. *Ann Meet NJ Mosq Exterm Assoc* 1938; 25: 176–185

- 138 Tawatsin A, Wratten SD, Scott RR, Thavara U, Techadamrongsin Y. Repellency of volatile oils from plants against three mosquito vectors. *J Vector Ecol* 2001; 26: 76–82
- 139 Bindra RL, Agarwal KK, Shukla YN, Agarwal SK, Kumar S. Essential oil formulations as mosquito repellents in floor cleaners. *Indian Perfumer* 2001; 45: 269–273
- 140 Thorsell W, Mikiver A, Malander I, Tunon H. Efficacy of plant extracts and oils as mosquito repellents. *Phytomedicine* 1998; 5: 311–323
- 141 Makhaik M, Naik SN, Tewary DK. Evaluation of anti-mosquito properties of essential oils. *J Sci Indian Res (India)* 2005; 64: 129–133
- 142 Yang P, Ma Y. Repellent effect of plant essential oils against *Aedes albopictus*. *J Vector Ecol* 2005; 30: 231–234
- 143 Moore SJ, Lenglet A, Hill N. Field evaluation of three plant-based insect repellents against malaria vectors in Vaca Diez Province, the Bolivian Amazon. *J Am Mosq Control Assoc* 2002; 18: 107–110
- 144 Kim S-I, Chang K-S, Yang Y-C, Kim B-S, Ahn Y-J. Repellency of aerosol and cream products containing fennel oil to mosquitoes under laboratory and field conditions. *Pest Manag Sci* 2004; 60: 1125–1130
- 145 Kim D-H, Kim S-I, Chang K-S, Ahn Y-J. Repellent activity of constituents identified in *Foeniculum vulgare* fruit against *Aedes aegypti* (Diptera: Culicidae). *J Agric Food Chem* 2002; 50: 6993–6996
- 146 Bhuyan M, Saxena N, Rao KM. Repellent property of oil fraction of garlic, *Allium sativum* (Linn.). *Indian J Exp Biol* 1974; 12: 575–576
- 147 Qualls WA, Xue R-D. Field evaluation of three botanical repellents against *Psorophora ferox*, *Aedes atlanticus*, and *Aedes mitchellae*. *J Am Mosq Control Assoc* 2009; 25: 379–381
- 148 Osmani Z, Anees I, Naidu MB. Insect repellent creams from essential oils. *Pesticides* 1972; 6: 19–21
- 149 Matsumoto A, Matsumoto T, Yamashita M, Nitsuta I, Kobayashi M, Mesaki J, Kashiwabara T. Harmful insect repellents containing edulan derivatives. Japanese Patent 05221808; 1993
- 150 Kobayashi A, Mizutani T. Harmful insect-repellent foot bands and method for repelling harmful insects using them. Japanese Patent 2004051564; 2004
- 151 Kang S-H, Kim M-K, Seo D-K, Noh D-J, Yang J-O, Yoon C, Kim G-H. Comparative repellency of essential oils against *Cx. pipiens pallens* (Diptera: Culicidae). *J Korean Soc Appl Biol Chem* 2009; 52: 353–359
- 152 Oshaghi MA, Ghalandari R, Vatandoost H, Shayeghi M, Kamalinejad M, Tourabi-Khaledi H, Abolhassani M, Hashemzadeh M. Repellent effect of extracts and essential oils of *Citrus limon* (Rutaceae) and *Melissa officinalis* (Labiatae) against main malaria vector, *Anopheles stephensi* (Diptera: Culicidae). *Iran J Public Health* 2003; 32: 47–52
- 153 Moore SJ, Darling ST, Sihuinchu M, Padilla N, Devine GJ. A low-cost repellent for malaria vectors in the Americas: results of two field trials in Guatemala and Peru. *Malar J* 2007; 6: 1–6
- 154 Maguranyi SK, Webb CE, Mansfield S, Russell RC. Are commercially available essential oils from Australian native plants repellent to mosquitoes? *J Am Mosq Control Assoc* 2009; 25: 292–300
- 155 Molyneux WM. Mosquito and/or flea control with *Leptospermum liversidgei* and its essential oil. Canadian Patent 2234476; 1999
- 156 Tawatsin A, Asavadachanukorn P, Thavara U, Wongsinkongman P, Bansidhi J, Boonruad T, Chavalittumrong P, Soonthornchareonnon N, Komalamisra N, Mulla MS. Repellency of essential oils extracted from plants in Thailand against four mosquito vectors (Diptera: Culicidae) and oviposition deterrent effects against *Aedes aegypti* (Diptera: Culicidae). *Southeast Asian J Trop Med Pub Health* 2006; 37: 915–931
- 157 Ding D, Sun H. Structural elucidation of an insect repellent in the essential oil of *Mentha haplocalyx* Briq. *Zhiwu Xuebao* 1983; 25: 62–66
- 158 Dubey VS. An overview of geraniol biosynthesis in palmarosa (*Cymbopogon martinii* Roxb. var. *motia*) and its biological and pharmacological properties. *Recent Progr Med Plants* 2006; 11: 75–97
- 159 Trongtokit Y, Rongsriyam Y, Komalamisra N, Apiwathnasorn C. Comparative repellency of 38 essential oils against mosquito bites. *Phytother Res* 2005; 19: 303–309
- 160 Ansari MA, Vasudevan P, Tandon M, Razdan RK. Larvicidal and mosquito repellent action of peppermint (*Mentha piperita*) oil. *Bioresour Technol* 2000; 71: 267–271
- 161 Tripathi AK, Prajapati V, Ahmad A, Aggarwal KK, Khanuja SPS. Piperitone oxide as toxic, repellent, and reproduction retardant toward malarial vector *Anopheles stephensi* (Diptera: Anophelinae). *J Med Entomol* 2004; 41: 691–698
- 162 Pavela R. Larvicidal property of essential oils against *Cx. quinquefasciatus* Say (Diptera: Culicidae). *Indian Crop Prod* 2009; 30: 311–315
- 163 Omolo MO, Okinyo D, Ndiege IO, Lwande W, Hassanali A. Repellency of essential oils of some Kenyan plants against *Anopheles gambiae*. *Phytochemistry* 2004; 65: 2797–2802
- 164 Dorfling EA, Mouton I. Insect repellents obtained from *Lippia* comprising monoterpenes. South African Patent ZA 9505853; 1997
- 165 Tripathi AK, Prajapati V, Aggarwal KK, Khanuja SP, Kumar S. Repellency and toxicity of oil from *Artemisia annua* to certain stored-product beetles. *J Econ Entomol* 2000; 93: 43–47
- 166 Das NG, Baruah I, Talukdar PK, Das SC. Evaluation of botanicals as repellents against mosquitoes. *J Vector Borne Dis* 2003; 40: 49–53
- 167 Xue R-D, Alp A, Barnard DR. Laboratory evaluation of toxicity of 16 insect repellents in aerosol sprays to adult mosquitoes. *J Am Mosq Control Assoc* 2003; 19: 271–274
- 168 Vatandoost H, Hanafi-Bojd AA. Laboratory evaluation of 3 repellents against *Anopheles stephensi* in the Islamic Republic of Iran. *East Mediterr Health J* 2008; 14: 260–267
- 169 Barnard DR, Xue R-D. Laboratory evaluation of mosquito repellents against *Aedes albopictus*, *Cx. nigripalpus*, and *Ochlerotatus triseriatus* (Diptera: Culicidae). *J Med Entomol* 2004; 41: 726–730
- 170 Sharma VP, Ansari MA, Razdan RK. Mosquito repellent action of neem (*Azadirachta indica*) oil. *J Am Mosq Control Assoc* 1993; 9: 359–360
- 171 Sharma SK, Dua VK, Sharma VP. Field studies on the mosquito repellent action of neem oil. *Southeast Asian J Trop Med Public Health* 1995; 26: 180–182
- 172 Dua VK, Nagpal BN, Sharma VP. Repellent action of neem cream against mosquitoes. *Indian J Malariol* 1995; 32: 47–53
- 173 Valecha N, Ansari MA, Prabhu S, Razdan RK. Preliminary evaluation of safety aspects of neem oil in kerosene lamp. *Indian J Malariol* 1996; 33: 139–143
- 174 Sharma VP, Ansari MA. Personal protection from mosquitoes (Diptera: Culicidae) by burning neem oil in kerosene. *J Med Entomol* 1994; 31: 505–507
- 175 Mishra AK, Singh N, Sharma VP. Use of neem oil as a mosquito repellent in tribal villages of Mandla District, Madhya Pradesh. *Indian J Malariol* 1995; 32: 99–103
- 176 Fradin MS, Day JF. Comparative efficacy of insect repellents against mosquito bites. *New Engl J Med* 2002; 347: 13–18
- 177 Ritchie SA, Williams CR, Montgomery BL. Field evaluation of new mountain sandalwood mosquito sticks and new mountain sandalwood botanical repellent against mosquitoes in North Queensland, Australia. *J Am Mosq Control Assoc* 2006; 22: 158–160
- 178 Sultana S, Ali M, Ansari SH, Bagri P. The effect of physical factors on chemical composition of the essential oil of *Eucalyptus citriodora* Hook. leaves. *J Essent Oilbearing Plants* 2008; 11: 69–74
- 179 Noosidum A, Prabaripai A, Chareonviriyaphap T, Chandrapatya A. Excito-repellency properties of essential oils from *Melaleuca leucadendron* L., *Litsea cubeba* (Lour.) Persoon, and *Litsea salicifolia* (Nees) on *Aedes aegypti* (L.) mosquitoes. *J Vector Ecol* 2008; 33: 305–312
- 180 Choochote W, Chaithong U, Kamsuk K, Jitpakdi A, Tippawangkosol P, Tuetun B, Champakaew D, Pitasawat B. Repellent activity of selected essential oils against *Aedes aegypti*. *Fitoterapia* 2007; 78: 359–364
- 181 Paluch G, Justingrodnitzky J, Bartholomay L, Coats J. Quantitative structure-activity relationship of botanical sesquiterpenes: spatial and contact repellency to the yellow fever mosquito, *Aedes aegypti*. *J Agric Food Chem* 2009; 57: 7618–7625
- 182 Hwang YS, Wu KH, Kumamoto J, Axelrod H, Mulla MS. Isolation and identification of mosquito repellents in *Artemisia vulgaris*. *J Chem Ecol* 1985; 11: 1297–1306
- 183 Odalo JO, Omolo MO, Malebo H, Angira J, Njeru PM, Ndiege IO, Hassanali A. Repellency of essential oils of some plants from the Kenyan coast against *Anopheles gambiae*. *Acta Trop* 2005; 95: 210–218
- 184 Chang K-S, Tak J-H, Kim S-I, Lee W-J, Ahn Y-J. Repellency of *Cinnamomum cassia* bark compounds and cream containing cassia oil to *Aedes aegypti* (Diptera: Culicidae) under laboratory and indoor conditions. *Pest Manag Sci* 2006; 62: 1032–1038
- 185 Traboulsi AF, El-Haj S, Tueni M, Taoubi K, Nader NA, Mrad A. Repellency and toxicity of aromatic plant extracts against the mosquito *Cx. pipiens molestus* (Diptera: Culicidae). *Pest Manag Sci* 2005; 61: 597–604
- 186 Beboukian RH, Weldon PJ. Spatial inhibitors, deterrents and repellents for mosquitoes and midges. World Patent 2007025197; 2007
- 187 Sugimoto M. Mosquito repellents of plant origin. Japanese Patent 2004210756; 2004

- 188 Hao H, Wei J, Dai J, Dui J. Host-seeking and blood-feeding behavior of *Aedes albopictus* (Diptera: Culicidae) exposed to vapors of geraniol, citral, citronellal, eugenol, or anisaldehyde. *J Med Entomol* 2008; 45: 533–539
- 189 Gu H-J, Cheng S-S, Lin C-Y, Huang C-G, Chen W-J, Chang S-T. Repellency of essential oils of *Cryptomeria japonica* (Pinaceae) against adults of the mosquitoes *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae). *J Agric Food Chem* 2009; 57: 11 127–11 133
- 190 Coats JR, Peterson CJ, Zhu J, Baker TC, Nemetz LT. Biorational terpenoid arthropod repellents obtained from plants. US Patent 6524605; 2003
- 191 Gladkikh SG. Repellents for arthropod carriers of infectious diseases. *Zhurnal Mikrobiologii Epidemiologii i Immunobiologii* 1953; 11: 61–65
- 192 Blagoveshchenskii DI, Bregetova NG, Monchadskii AS. New deterrent substances for protecting man against the attacks of mosquitoes. *Bull Acad Sci USSR, Div Chem Sci (Engl. Transl.)* 1943; 40: 119–122
- 193 Cantrell CL, Klun JA, Bryson CT, Kobaisy M, Duke SO. Isolation and identification of mosquito bite deterrent terpenoids from leaves of American (*Callicarpa americana*) and Japanese (*Callicarpa japonica*) beautyberry. *J Agric Food Chem* 2005; 53: 5948–5953
- 194 Kim SI, Yeon SH, Yoon HS, Ahn YJ. Natural repellent containing essential oils derived from *Zanthoxylum armatum* and/or *Zanthoxylum piperitum*. Korean Patent 955344; 2010
- 195 Kim GH, Kang SH. Mosquito-repellent composition. Korean Patent 756397; 2007
- 196 Darling ST. Mosquito repellent composition for skin. World Patent 2007082306; 2007
- 197 Georges PS. Material controllingly releasing a liquid active substance by evaporation and a mosquito repellent bracelet comprising said material. World Patent 2007026082; 2007
- 198 Thanispong K, Achee NL, Grieco JP, Bangs MJ, Suwonkerd W, Prabaripai A, Chauhan KR, Chareonviriyaphap R. A high throughput screening system for determining the three actions of insecticides against *Aedes aegypti* (Diptera: Culicidae) populations in Thailand. *J Med Entomol* 2010; 47: 833–841
- 199 Hwang YS, Schultz GW, Mulla MS. Structure-activity relationships of unsaturated fatty acids as mosquito ovipositional repellents. *J Chem Ecol* 1984; 10: 145–151
- 200 Nataraj S. Mosquito repellent comprising natural oils, turmeric powder, and camphor. Indian Patent 2003MA00347; 2005
- 201 Lee YI. Natural vegetable essential oil composition having sanitary insect pest exterminating and antibacterial effects and formulation thereof. Korean Patent 2000072252; 2000
- 202 Gries R, Campbell C, Khaskin G, Avelino N, Gries G. Allyl sulfide compounds, and compositions and methods using said compounds for repelling blood-feeding arthropods. World Patent 2008116321; 2008
- 203 Hallahan DL, Manzer LE. Preparation of dihydronepetalactone as insect repellent. International Patent 2005034626; 2005
- 204 Ravid U, Putievsky E, Katzir I, Ikan R. Determination of the enantiomeric composition of terpinen-4-ol in essential oils using a permethylated β -cyclodextrin coated chiral capillary column. *Flav Fragr J* 1992; 7: 49–52