



Mathematisches
Forschungsinstitut
Oberwolfach

Member of the



Oberwolfach Preprints

OWP 2020 - 13

BERRY GROISMAN, MICHAEL MC GETTRICK,
MEHDI MHALLA AND MARCIN PAWLOWSKI

How Quantum Information Can Improve Social
Welfare

Mathematisches Forschungsinstitut Oberwolfach gGmbH
Oberwolfach Preprints (OWP) ISSN 1864-7596

Oberwolfach Preprints (OWP)

The MFO publishes a preprint series **Oberwolfach Preprints (OWP)**, ISSN 1864-7596, which mainly contains research results related to a longer stay in Oberwolfach, as a documentation of the research work done at the MFO. In particular, this concerns the Research in Pairs-Programme (RiP) and the Oberwolfach-Leibniz-Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it on its website as well as by hard copy. Every RiP group or Oberwolfach-Leibniz-Fellow may receive on request 30 free hard copies (DIN A4, black and white copy) by surface mail.

The full copyright is left to the authors. With the submission of a manuscript, the authors warrant that they are the creators of the work, including all graphics. The authors grant the MFO a perpetual, non-exclusive right to publish it on the MFO's institutional repository.

In case of interest, please send a **pdf file** of your preprint by email to rip@mfo.de or owlf@mfo.de, respectively. The file should be sent to the MFO within 12 months after your stay as RiP or OWLF at the MFO.

There are no requirements for the format of the preprint, except that the introduction should contain a short appreciation and that the paper size (respectively format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX). Additionally, each preprint will get a Digital Object Identifier (DOI).

We cordially invite the researchers within the RiP or OWLF programme to make use of this offer and would like to thank you in advance for your cooperation.

Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)
Schwarzwaldstrasse 9-11
77709 Oberwolfach-Walke
Germany

Tel +49 7834 979 50
Fax +49 7834 979 55
Email admin@mfo.de
URL www.mfo.de

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO.
Copyright of the content is held by the authors.

DOI 10.14760/OWP-2020-13

How Quantum Information can improve Social Welfare

Berry Groisman¹, Michael Mc Gettrick², Mehdi Mhalla³, and Marcin Pawłowski⁴

¹Department of Applied Mathematics and Theoretical Physics, University of Cambridge, UK; email: b.groisman@damtp.cam.ac.uk

²School of Mathematics, Statistics and Applied Mathematics, National University of Ireland, Galway, Ireland; email:

michael.mcgettrick@nuigalway.ie

³Université Grenoble Alpes, Grenoble, France; email:

mhallam@univ-grenoble-alpes.fr

⁴International Centre for Theory of Quantum Technologies, University of Gdansk, Wita Stwosza 63, 80-308 Gdansk, Poland; email:

dokmpa@univ.gda.pl

July 17, 2020

Abstract

In [2, 18, 5, 19, 4] it has been shown that quantum resources can allow us to achieve a family of equilibria that can have sometimes a better social welfare, while guaranteeing privacy. We use graph games to propose a way to build non-cooperative games from graph states, and we show how to achieve an unlimited improvement with quantum advice compared to classical advice.

1 Introduction

An important tool in analysing games is the concept of *Nash equilibrium* [17], which represents situations where no player has incentive to deviate from their strategy. This corresponds to situations observed in real life, with applications in economics, sociology, international relations, biology, etc. All equilibria do not have the same

social welfare, i.e. the average payoff is different from one equilibrium to another. Games of incomplete information can exhibit better equilibria if players use a resource – a general correlation, Q . Such correlation can be viewed as a resource produced by a mediator to give *advice* to the players. The concept of advice generalizes the notion of Nash equilibrium to a broader class of equilibria [3]. All such equilibria can be classified according to the properties of the resource correlation. Three classes can be identified in addition to Nash equilibria (no correlation), namely general communication equilibria (Comm) [9], where Q is unrestricted, belief-invariant equilibria (BI) [10, 11, 14, 15] and correlated equilibria (Corr) [3]. The canonical versions of these equilibria form a sequence of nested sets within the set of canonical correlations:

$$\text{Nash} \subset \text{Corr} \subset \text{BI} \subset \text{Comm}.$$

It was demonstrated that there exist games where BI equilibria can outperform Corr equilibria [18] (in terms of a social welfare (SW) of a game) as well as games where BI equilibria outperform any non-BI equilibria. In [19] the work of [18] is extended into the quantum domain.

Auletta et al. [2] introduce quantum correlated equilibria as a subclass of BI equilibria and show that quantum correlations can achieve optimal SW. This provides the link with quantum nonlocality, where quantum resources are used to produce *non-signalling* correlations. In this context, belief invariance describes the largest class of correlations that obey *relativistic causality*. The role of *quantum entanglement* as quantum-social welfare advice was further studied in [4].

A characteristic feature of belief-invariance is that it ensures privacy – the other players involved in the game have no information about the input one player sent to the resource.

To obtain the canonical form of the games, [16] show that one can suppose that the output of the correlation resource is the answer the players give by delegating the extra computation (from game question to input to the box and from output of the box to players' answer) to the mediator. Therefore, quantum equilibria can be

reached in a setting where players each measure quantum systems or, equivalently, by just having a central system providing advices by measuring a quantum device.

Ref. [2] highlights several open questions. In particular,

- (1) Whether any full-coordination game (a.k.a. a *non-local game* in quantum physics and computer science communities) can be converted into a conflict-of-interests game. Ref. [18] gives an example of a two-player variant of the CHSH game, while [2] extends their result to an n -player game in which there exists a BI equilibrium which is better than any Corr equilibrium.
- (2) How can we get a large separation between the expected payoff for the quantum and correlated equilibrium cases, and what is the upper bound for the separation? In the case of two-player full coordination games this question was settled in [7, 13]. Are there conflict-of-interest games which exhibit large separation?

In this paper, we provide a natural way to convert graph games (and more generally stabiliser games) into conflict-of-interest games, and we show how we can create unbounded separation by increasing the number of players or using penalty techniques (a negative payoff).

An interesting feature in these games compared to the usual pseudo-telepathy scenarios studied in quantum information is the notion of *involvement* [1, 16], which allows one to define some interesting scenarios in non-cooperative games and which exhibits novel features, e.g. unlimited separation. If a player participates in the game but is not involved (on a particular round) it means that their strategy is not taken into account when determining the win/lose outcome. However, they do receive a corresponding payoff.

Using these games one can build games with bounded personal utilities v_0, v_1 on $O(\log(\frac{1}{\epsilon}))$ players ensuring $\frac{CSW(G)}{QSW(G)} \leq \epsilon$, where CSW/QSW are the Classical/Quantum Social Welfares, respectively.

The paper is organized as follows. In Sec. 2 we describe graph games which are the underlying non-local games used to define our games. In Sec. 3 we define a non-collaborative game as a modification of the collaborative games by introducing unequal payoffs corresponding to answers 0 and 1 of each player, and discuss

the corresponding quantum perfect strategy. We consider a particular version of graph games from the cycle on five vertices. Sec. 4 discusses variations of non-collaborative games based on the cycle on five vertices. Finally, Sec. 5 shows how one can amplify the quantum advantage by adding a penalty for wrong answers and by increasing the number of players.

2 Graph games

Non-local games play a key role in Quantum Information theory. They can be viewed as a setting in which players that are not allowed to communicate receive some inputs and have to produce some outputs, and there is a winning/losing condition depending globally on their outputs for each input. Particular types of games are pseudo-telepathy games [6] which are games that can be won perfectly using quantum resources but that are impossible to win perfectly without communication when the players have access only to shared randomness. Multipartite collaborative games ($MCG(G)$) are a family of pseudo-telepathy games based on certain types of quantum states called *graph states*. The players are identified with vertices of the graph and have a binary input/output each with the winning/losing conditions built using the stabilisers of the graph states.

The combinatorial game¹ MCG with n players consists in asking the players questions: for each question q , each player i receives one bit q_i as input and answers one bit a_i . They can either all win or all lose depending on their answer, with winning/losing conditions described by a set $\{(q, I(q), b(q))\}$ where

- $q \in \{0, 1\}^n$ is a valid question in which each player i gets the bit q_i and in the subgraph of the vertices corresponding to players receiving one, all vertices have even degree. Let $I_1 = \{i, q_i = 1\}$ and $G' = G_{|I_1}$, a question is valid if each vertex of G' has an even number of neighbors in G'
- $I(q) \subset [n]$ is a subset of players that are called '*involved*' in the question as the sum (modulo 2) of their answers determines the winning/losing condition according to the bit $b(q)$:

¹without considering probability distributions

- $b(q)$ is defined such that the players win the game when the question is q if the sum of the answers of the involved players is equal to the parity of the number of edges of the subgraph of the vertices corresponding to players receiving one:

$$\sum_{i \in I(q)} a_i = b(q) = |E(G')| \bmod 2.$$

For instance the game associated to the cycle on 5 elements $MCG(C_5)$ is defined by

- When the question is $q = 11111$ (each player has input 1), the players lose if the binary sum of their answer is 0, *i.e.* $\sum_{i=0}^4 a_i = 0 \bmod 2$, and win otherwise.
- When the question contains 010 for three players corresponding to three adjacent vertices, the players lose if the binary sum of the answer of these three players is 1 *i.e.* $a_{i-1} + a_i + a_{i+1} = 0 \bmod 2$ when q contains $0_{i-1}1_i0_{i+1}$.
- The players win otherwise.

A variation of this game can be done by reducing the set of valid questions, for instance in the above set-up the questions of the second type have only three players “involved”, so a first version could be to chose only 5 questions of the second type and give always 0 as advice to the non-involved players. This is the game studied as an example in [16].

An important point is that the notion of involvement in MCG games is absent in unique games and introduces situations where the players might change their strategy (answer) without changing the winning/losing status of the global strategy.

To analyse these games and the strategies, one can imagine a scenario where there is one special player representing Nature who is playing against the other players. The strategy of Nature is therefore a probability distribution over the questions that we study here (as is standard in game theory) as a known function on the set of questions $w : T \rightarrow [0, 1]$ such that $\sum_{t \in T} w(t) = 1$. The games will be therefore defined by equipping the combinatorial game with a probability distribution over the questions.

3 Defining non-collaborative games

Like in multipartite collaborative graph games $MCG(G)$, we associate a non-collaborative game $NC(G)$ to each graph. We differentiate the payoff of the players using the value of their output: If the global answer wins in the non-local game, each player gets v_1 if they answer 1 and v_0 if they answer 0. If the global answer loses, they get 0.

To match the traditional terminology used in game theory the output from now on will be called *strategy*, and the input called *type*. The payoff is called *utility* and the social welfare is the average of the utilities over the players.

A non-collaborative game $NC(G)$ is thus defined from $MCG(G)$ as follows

- The considered types are $T \subset \{0, 1\}^n$ where n is the number of vertices of G .
- As in MCG , to each type $t \in T$ corresponds an associated involved set $I(t)$ of players, and an expected binary answer $b(t)$.
- As in MCG , the losing set is

$$\mathcal{L} = \{(s, t), \sum_{i \in I(t)} s_i \neq b(t) \pmod{2}\}.$$

We say that the players using a strategy s , given a type t , collectively win the game when the sum of the local strategies of the involved players is equal to the requested binary answer modulo 2.

- the payoff function is:

$$u_j(s|t) = \begin{cases} v_{s_j} & \text{if } (s, t) \notin \mathcal{L} \\ 0 & \text{Otherwise} \end{cases}$$

Firstly we consider the cycle on five vertices C_5 . We define $NC_{00}(C_5)$ based on the non-local game $MCG(C_5)$ studied in [1, 16]. For questions which involve three players, both non-involved players have type 0 (see Figure 1).

We consider the game with the type probability distribution $w(t) = 1/6$ for all the types.

The quantum perfect strategy for $NC(G)$ is obtained when the players each




Figure 1: $NC_{00}(C_5)$: Square nodes indicate a 1 in the associated type, while circular nodes indicate a 0. Involved players in each case are shaded in **red**.

| Type | Involved set | Binary answer |
|---------------|------------------------------|---------------|
| $T_a = 11111$ | $I(T_a) = \{0, 1, 2, 3, 4\}$ | $b(T_0) = 1$ |
| $T_0 = 10000$ | $I(T_0) = \{0, 1, 4\}$ | $b(T_0) = 0$ |
| $T_1 = 01000$ | $I(T_1) = \{0, 1, 2\}$ | $b(T_1) = 0$ |
| $T_2 = 00100$ | $I(T_2) = \{1, 2, 3\}$ | $b(T_2) = 0$ |
| $T_3 = 00010$ | $I(T_3) = \{2, 3, 4\}$ | $b(T_3) = 0$ |
| $T_4 = 00001$ | $I(T_4) = \{3, 4, 0\}$ | $b(T_4) = 0$ |

Table 1: $NC_{00}(G)$ game.

have a qubit from graph state $|G\rangle$ [1]. Each player i measures their qubit according to their type t_i , getting a quantum advice representing their part of the quantum strategy s_i [1]. From the study of $MCG(G)$ we have

Theorem 1. *If all the players collaborate (follow the quantum advice) then for any probability distribution over the types, the utility of each player is $(v_0 + v_1)/2$.*

Proof. The output of each quantum measurement provides uniformly all the possible answers. □

3.1 Is the quantum pseudo-telepathy solution a Nash equilibrium?

As the players now have an incentive to answer 1, they can sacrifice always getting a good answer to maximize their utility. Indeed, in the previous game, each player is always involved when they get type 1 and with probability 1/2 when they get type 0; getting the wrong answer in that case only costs v_0 .

Without loss of generality we consider $v_1 \geq v_0$. The players now have an incentive to answer 1, because they might be able to maximize their utility by allowing the non-zero probability of a wrong answer. Indeed, in the previous game, $NC_{00}(C_5)$, if the player gets type 1 then they are certain that they are involved, and they won't gain by defecting (not following advice). However, if their type is 0, then the probability of them being involved is 1/2, and so there is a fifty percent chance that they will benefit from always answering 1 while not compromising the winning combination. Getting the wrong answer in that case only costs v_0 .

Theorem 2. *Let $p_{inv}^{(i)}(t_i, s_i)$ be the probability for the player i who gets type t_i and advice s_i to be involved*

Then, in $NC(G)$, the quantum advice gives a belief-invariant Nash equilibrium iff

$$\frac{v_0}{v_1} \geq (1 - p),$$

where

$$p = \min_i \min_{t_i} p_{inv}^{(i)}(t_i, 0).$$

Proof. If the advice is $s_i = 1$ then the winning payoff is already v_1 . Consider the case when player i is given the advice $s_i = 0$ (which would lead to payoff v_0 in the winning case). If the player defects then the difference of utility is $-v_0 p_{inv}^{(i)}(t_i, 0) + (1 - p_{inv}^{(i)}(t_i, 0))(v_1 - v_0)$. So the strategy is a Nash-equilibrium when $(1 - p_{inv}^{(i)}(t_i, 0))v_1 \leq v_0$, i.e $v_0/v_1 \geq 1 - p_{inv}^{(i)}(t_i, 0)$. This inequality has to hold for all types and all players.

□

For $NC_{00}(C_5)$, $p_{inv}^{(i)}(0, 0) = 1/2$ and therefore the quantum nonlocal strategy is

an equilibrium only when $v_0/v_1 \geq 1/2$.

One important characteristic of an equilibrium is the *Social Welfare*, which is the average utility of the players.

As a direct consequence of Theorem 1 the average social welfare of the quantum strategy is independent on the graph

$$QSW(NC(G)) = \frac{v_0 + v_1}{2}.$$

Note that the non collaborative games defined have a special feature that we call *guaranteed value*: in any run of the game players following the quantum strategy receive their expected payoff with probability 1.

4 Some versions of $NC(C_5)$

In this section we study the game $NC_{00}(C_5)$ and then introduce a number of modifications in order to improve the quantum advantage (ratio of quantum social welfare to correlated social welfare) and also to symmetrize the game such that the players get 0 and 1 with same probability or have the same probability of being involved regardless of whether their type is 0 or 1.

4.1 Study of $NC_{00}(C_5)$

Pure Nash equilibria can be described by local functions: each player having one local type bit and one strategy bit to produce, can locally act as follows:

- $0 \rightarrow 0, 1 \rightarrow 0$ constant function 0 denoted **0**
- $0 \rightarrow 1, 1 \rightarrow 1$ constant function 1 denoted **1**
- $0 \rightarrow 0, 1 \rightarrow 1$ Identity function denoted **2**
- $0 \rightarrow 1, 1 \rightarrow 0$ NOT function denoted **3**

The set of pure Nash equilibria depends on the ratio v_0/v_1 . There are 20/25/40 pure Nash equilibria (4/4/6 up to symmetry) when v_0/v_1 lies within the interval $[0, 1/3]$, $[1/3, 1/2]$ or $[1/2, 1]$ respectively (see Table 2).

| Local functions | Players utility [$\times 6$] | | | | | $SW[\times 30]$ |
|-----------------------------|--------------------------------|---------------|---------------|---------------|---------------|-----------------|
| $v_0/v_1 \leq 1/3$ | | | | | | |
| 2 1 1 1 1 | $2v_0 + v_1$ | $3v_1$ | $3v_1$ | $3v_1$ | $3v_1$ | $2v_0 + 13v_1$ |
| 3 3 1 1 1 | $2v_0 + v_1$ | $2v_0 + v_1$ | $3v_1$ | $3v_1$ | $3v_1$ | $v_0 + 11v_1$ |
| 3 1 3 1 1 | $2v_0 + v_1$ | $3v_1$ | $2v_0 + v_1$ | $3v_1$ | $3v_1$ | $4v_0 + 11v_1$ |
| 3 3 3 3 1 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| $1/3 \leq v_0/v_1 \leq 1/2$ | | | | | | |
| 1 3 1 1 0 | $5v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $7v_0 + 18v_1$ |
| 2 2 1 1 1 | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_1$ | $6v_0 + 19v_1$ |
| 3 3 1 1 1 | $2v_0 + v_1$ | $2v_0 + v_1$ | $3v_1$ | $3v_1$ | $3v_1$ | $4v_0 + 11v_1$ |
| 3 3 3 3 1 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| $v_0/v_1 \geq 1/2$ | | | | | | |
| 3 2 1 1 0 | $2v_0 + 3v_1$ | $4v_0 + v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $11v_0 + 14v_1$ |
| 1 3 1 1 0 | $5v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $7v_0 + 18v_1$ |
| 2 2 1 1 1 | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_1$ | $6v_0 + 19v_1$ |
| 3 3 1 2 1 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $4v_0 + v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| 3 3 3 3 1 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| 3 2 3 2 2 | $2v_0 + 3v_1$ | $4v_0 + v_1$ | $2v_0 + 3v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $14v_0 + 11v_1$ |

Table 2: Nash equilibria for three intervals of the value v_0/v_1 . Note that the critical values $1/2$ and $1/3$ have union of both tables as equilibria.

We can see that most of these equilibria (all of them when $v_0/v_1 \geq 1/2$) correspond to local functions winning for the 5 types.

When $v_0 = 2/3$ and $v_1 = 1$ then the quantum social welfare of the pseudotelepathy strategy is $QSW = 0.83$ whereas the best classical social welfare $CSW = 0.77$.

As noted in section 3.1 the probability of being involved in NC_{00} is $p(1, s) = 1$ and $p(0, s) = 1/2$ and the quantum pseudotelepathy measurements strategy is an

equilibrium if $v_0/v_1 \geq 1/2$.

Similar behavior can be seen with Pareto equilibria (ones in which local utility cannot improve without reducing the outcome of someone else): see Appendix.

Recall that the characteristic feature of $NC_{00}(C_5)$ is that each player has unequal probabilities of getting different types. The game can be symmetrized by changing the types of the non-involved players from 00 to 01, as shown in the next section.

4.2 Comments on $NC_{01}(C_5)$

We define a second variant from $MCG(C_5)$: $NC_{01}(C_5)$ where any player gets the types 0 and 1 with probability 1/2 by adding an extra 1 for a non-involved player in the types so that $T_i = 0_{i-1}1_i0_{i+1}1_{i+2}0_{i+3}$: see Table 3.

If the type probability distribution is $w(t) = 1/6$ for all the types, then one can see that any player is involved with probability 2/3 whether their input is 0 or 1, i.e. $p_{\text{inv}}^{(i)}(0, 0) = p_{\text{inv}}^{(i)}(1, 0) = 2/3$. Hence, by Theorem 2, the quantum strategy of MCG produces a Nash equilibrium iff $v_0/v_1 \geq 1/3$. Thus, one of the benefits of this variant is that quantum Nash equilibria exist at a lower ratio v_0/v_1 .

Note that in this version each player is getting a perfect random bit as advice : $p(a = 1) = p(a = 0) = 1/2$.

When $v_0 = 2/3$ and $v_1 = 1$ then the quantum social welfare of the pseudotelepathy strategy is $QSW = 0.83$ whereas the best classical social welfare is $CSW = 0.78$.

| Type | Involved set | Binary answer |
|---------------|------------------------------|---------------|
| $T_a = 11111$ | $I(T_a) = \{0, 1, 2, 3, 4\}$ | $b(T_a) = 1$ |
| $T_0 = 10100$ | $I(T_0) = \{0, 1, 4\}$ | $b(T_0) = 0$ |
| $T_1 = 01010$ | $I(T_1) = \{0, 1, 2\}$ | $b(T_1) = 0$ |
| $T_2 = 00101$ | $I(T_2) = \{1, 2, 3\}$ | $b(T_2) = 0$ |
| $T_3 = 10010$ | $I(T_3) = \{2, 3, 4\}$ | $b(T_3) = 0$ |
| $T_4 = 01001$ | $I(T_4) = \{3, 4, 0\}$ | $b(T_4) = 0$ |

Table 3: $NC_{01}(G)$ game (Here the players are identified with the integers modulo 5).

4.3 Comments on $NC_{00,0}(C_5)$

A modification of a different kind consists in adding more questions from the stabiliser. As the first example of this kind we define a game $NC_{00,0}(C_5)$, where the additional family of questions has four involved players with the non-involved player getting type 0, as specified by Table 4.

| Type | Involved set | Binary answer |
|--|---|------------------|
| $T_a = 11111$ | $I(T_a) = \{0, 1, 2, 3, 4\}$ | $b(T_0) = 1$ |
| $T_{i_1} = 0_{i_1-2}0_{i_1-1}1_{i_1}0_{i_1+1}0_{i_1+2}$ $i_1 \in \{0, \dots, 4\}$ | $I(T_{i_1}) = \{i_1 - 1, i_1, i_1 + 1\}$ | $b(T_{i_1}) = 0$ |
| $T_{i_2} = 0_{i_2-1}1_{i_2}0_{i_2+1}1_{i_2+2}0_{i_2+3}$ $i_2 \in \{0, \dots, 4\}$ | $I(T_{i_2}) = \{i_2 - 1, i_2, i_2 + 2, i_2 + 3\}$ | $b(T_{i_2}) = 0$ |

Table 4: $NC_{00,0}(G)$ game.

For $v_1 = 1$, $v_0 = \frac{2}{3}$, and the probability distribution $w(T_a) = \frac{3}{13}$, $w(T_{i_1}) = w(T_{i_2}) = \frac{1}{13}$ we get a CSW of 0.72 versus a QSW of 0.83.

Note that each player gets types 0 and 1 with different probabilities. In fact, it is simple to show that no choice of w_1, w_2 and w_3 can make these probabilities equal. However, it is possible to modify the set of types so that equality becomes possible, as shown in the following.

4.4 Comments on $NC_{00,01,0}(C_5)$

We increase the set of types using other questions from the stabiliser: We define a game $NC_{00,01,0}(C_5)$ for which with a suitable choice of probability distribution the players get 0 and 1 with the same probability.

We consider this game with type probability distributions given by $w(T_a) = 3/13$, $w(T_{i_1}) = 1/26$, $w(T_{i_2}) = 1/26$ and $w(T_{i_3}) = 1/13$.

The involvement probabilities satisfy $P_{inv}(1) > P_{inv}(0) = 8/13$ and the best Classical Social Welfare with $v_0 = 2/3$, $v_1 = 1$ is CSW = 0.72 versus a QSW of 0.83.

Note that even though the types T_{i_2} and T_{i_3} are similar, the involved sets and thus the utilities are different. However, if one wants to restrict to scenarios in

| Type | Involved set | Binary answer |
|--|---|------------------|
| $T_a = 11111$ | $I(T_a) = \{0, 1, 2, 3, 4\}$ | $b(T_0) = 1$ |
| $T_{i_1} = 0_{i_1-2}0_{i_1-1}1_{i_1}0_{i_1+1}0_{i_1+2}$ $i_1 \in \{0, \dots, 4\}$ | $I(T_{i_1}) = \{i_1 - 1, i_1, i_1 + 1\}$ | $b(T_{i_1}) = 0$ |
| $T_{i_2} = 0_{i_2-1}1_{i_2}0_{i_2+1}1_{i_2+2}0_{i_2+3}$ $i_2 \in \{0, \dots, 4\}$ | $I(T_{i_2}) = \{i_2 - 1, i_2, i_2 + 1\}$ | $b(T_{i_2}) = 0$ |
| $T_{i_3} = 0_{i_3-1}1_{i_3}0_{i_3+1}1_{i_3+2}0_{i_3+3}$ $i_3 \in \{0, \dots, 4\}$ | $I(T_{i_3}) = \{i_2 - 1, i_2, i_2 + 2, i_2 + 3\}$ | $b(T_{i_3}) = 0$ |

Table 5: $NC_{00,01,0}(G)$ game.

which the utility can be deterministically determined from the type, one can just add an extra player with a type allowing to distinguish the different cases and with utility the average utility of the other players independently of his/her action.

5 Quantum vs correlation separation

In [2] it is asked as an open question whether the separation between classical and quantum social welfare is bounded. We show in this section how two families of amplification techniques can increase the separation by adding a penalty for wrong answers and then by increasing the number of players.

5.1 Wrong answer penalty

A possible technique is to penalize bad answers more, using the fact that classical functions always produce a bad answer for some question. Instead of getting 0 when losing we generalize so that each player gets $-N_g v_1$ if they answer 1 and $-N_g v_0$ if they answer 0, where N_g can be seen as the penalty for giving a wrong answer. If $\delta_{(s,t),\mathcal{L}} = 1$ if $(s, t) \in \mathcal{L}$ and 0 otherwise, and N_g is a positive number, then

$$u_j(s|t) = (-N_g)^{\delta_{(s,t),\mathcal{L}}} v_{s_j}$$

For $NC_{01}(C_5)$ as soon as $N_g > 3v_1$ there exists only two classical Nash equilibria:

- All 0 with a social welfare of $\frac{-N_g v_0 + 5v_0}{6}$ and

- All NOT with a social welfare of $\frac{-N_g v_0 + 2v_0 + 3v_1}{6}$.

Therefore the classical social welfare decreases linearly with the penalty while the quantum average social welfare remains $\frac{v_1 + v_0}{2}$.

5.2 Distributed parallel repetition

The distributed parallel composition of nonlocal games appears in [12] for the study of non signaling correlations and also in [8] where it is called k -fold repetition. k groups of players play at the same time and they win collectively if all the groups win their game.

Theorem 3. *There exists games with bounded personal utilities v_0, v_1 on $O(\log(\frac{1}{\epsilon}))$ players ensuring $\frac{CSW(G)}{QSW(G)} \leq \epsilon$ for the ratio best classical social welfare over quantum social welfare with guaranteed value.*

Proof. It is easy to bound the utility in these settings as for any strategy in a repeated game. If a player p is involved in the strategy S_j but is not involved in the strategy S_i of another group then his utility is conditioned by the fact that the S_i strategy wins to receive a positive utility and

$$u^p(S_i \times S_j) \leq p_{win}(S_i)u^p(S_j)$$

As the quantum strategy obtained from following the nonlocal advice always wins, the QSW remains unchanged whereas the CSW decreases. For instance $CSW(k\text{-fold } NC_{00}(C_5)) = \frac{5^k}{6} CSW(NC_{00}(C_5))$.

Therefore using these games one can build games with bounded personal utilities v_0, v_1 on $O(\log(\frac{1}{\epsilon}))$ players ensuring $\frac{CSW(G)}{QSW(G)} \leq \epsilon$ □

6 Conclusion

We have used properties of multipartite graph games to define conflict of interest games, and shown that by combining such games the ratio classical social welfare / quantum social welfare can go to zero.

One can easily extend to stabilizer games [8] to have any number of types and possible strategies.

As pointed out by [2], quantum advice equilibria can be reached without needing a trusted mediator, furthermore they ensure privacy as they are belief invariant. Some other features may be emphasized if we define Nash equilibria using pseudotelepathy games: such situations ensure a guaranteed utility and they are also better when analysing the maximal minimal utility. It may be interesting to investigate further how this guaranteed value property for some quantum equilibria can be used. On the other hand it would also be interesting to investigate how relaxing the guaranteed win requirement might allow to increase the QSW even further.

The possibility of potentially unlimited improvement of social welfare while preserving belief invariance is therefore a strong motivation to consider classical payoff tables that arise for usual situations in which Nash equilibria occur and play an important role. For example, in routing problems an advice provider could use a quantum advice system as follows. To calculate the advice to send to each player, the advice provider should either (a) send a rotated qubit to each player (who will then measure their qubit to get the answer), or, in a trusted setting, (b) perform a quantum measurement and send a classical message.

Acknowledgement

This research was supported through the program “Research in Pairs” by the Mathematisches Forschungsinstitut Oberwolfach in 2019. The authors also acknowledge the “*Investissements d’avenir*” (ANR-15-IDEX-02) program of the French National Research Agency, NCN grant Sonata UMO-2014/14/E/ST2/00020 and thank Sidney Sussex College, Cambridge for support.

Appendix

Pareto equilibria for NC_{00} when $v_0/v_1 \leq 1/3$, Nb solutions : 121, Nb distinct

equilibria : 18

| Local functions | Players utility [$\times 6$] | | | | | $SW[\times 30]$ |
|-----------------|--------------------------------|---------------|---------------|---------------|---------------|-----------------|
| 2 1 1 0 0 | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $5v_0$ | $13v_0 + 12v_1$ |
| 3 3 2 0 0 | $v_0 + 2v_1$ | $v_0 + 2v_1$ | $v_0 + 2v_1$ | $3v_0$ | $3v_0$ | $9v_0 + 6v_1$ |
| 3 2 1 1 0 | $2v_0 + 3v_1$ | $4v_0 + v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $11v_0 + 14v_1$ |
| 1 3 1 1 0 | $5v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $7v_0 + 18v_1$ |
| 1 3 3 1 0 | $5v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $5v_1$ | $5v_0$ | $7v_0 + 18v_1$ |
| 2 3 1 2 0 | $3v_0 + 2v_1$ | $v_0 + 4v_1$ | $5v_1$ | $3v_0 + 2v_1$ | $5v_0$ | $12v_0 + 13v_1$ |
| 3 2 2 3 0 | $v_0 + 4v_1$ | $4v_0 + v_1$ | $4v_0 + v_1$ | $v_0 + 4v_1$ | $5v_0$ | $15v_0 + 10v_1$ |
| 2 1 1 1 1 | $2v_0 + v_1$ | $3v_1$ | $3v_1$ | $3v_1$ | $3v_1$ | $2v_0 + 13v_1$ |
| 2 2 1 1 1 | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_1$ | $6v_0 + 19v_1$ |
| 3 3 1 1 1 | $2v_0 + v_1$ | $2v_0 + v_1$ | $3v_1$ | $3v_1$ | $3v_1$ | $4v_0 + 11v_1$ |
| 3 1 3 1 1 | $2v_0 + v_1$ | $3v_1$ | $2v_0 + v_1$ | $3v_1$ | $3v_1$ | $4v_0 + 11v_1$ |
| 3 3 1 2 1 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $4v_0 + v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| 3 3 2 2 1 | $2v_0 + 3v_1$ | $v_0 + 4v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $9v_0 + 16v_1$ |
| 3 2 3 2 1 | $v_0 + 2v_1$ | $2v_0 + v_1$ | $2v_0 + v_1$ | $2v_0 + v_1$ | $3v_1$ | $7v_0 + 8v_1$ |
| 3 2 2 3 1 | $v_0 + 2v_1$ | $v_0 + 2v_1$ | $v_0 + 2v_1$ | $v_0 + 2v_1$ | $3v_1$ | $4v_0 + 11v_1$ |
| 3 3 3 3 1 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| 3 2 3 2 2 | $2v_0 + 3v_1$ | $4v_0 + v_1$ | $2v_0 + 3v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $14v_0 + 11v_1$ |
| 3 3 3 3 3 | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $5v_0 + 20v_1$ |

Pareto equilibria for NC_{00} when $1/3 \leq v_0/v_1 \leq 1/2$ Nb solutions : 91, Nb distinct

equilibria : 14

| Local functions | Players utility [$\times 6$] | | | | | $SW[\times 30]$ |
|-----------------|--------------------------------|---------------|---------------|---------------|---------------|-----------------|
| 2 1 1 0 0 | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $5v_0$ | $13v_0 + 12v_1$ |
| 3 2 1 1 0 | $2v_0 + 3v_1$ | $4v_0 + v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $11v_0 + 14v_1$ |
| 1 3 1 1 0 | $5v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $7v_0 + 18v_1$ |
| 1 3 3 1 0 | $5v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $5v_1$ | $5v_0$ | $7v_0 + 18v_1$ |
| 2 3 1 2 0 | $3v_0 + 2v_1$ | $v_0 + 4v_1$ | $5v_1$ | $3v_0 + 2v_1$ | $5v_0$ | $12v_0 + 13v_1$ |
| 3 2 2 3 0 | $v_0 + 4v_1$ | $4v_0 + v_1$ | $4v_0 + v_1$ | $v_0 + 4v_1$ | $5v_0$ | $15v_0 + 10v_1$ |
| 2 2 1 1 1 | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_1$ | $6v_0 + 19v_1$ |
| 3 3 1 1 1 | $2v_0 + v_1$ | $2v_0 + v_1$ | $3v_1$ | $3v_1$ | $3v_1$ | $4v_0 + 11v_1$ |
| 3 3 1 2 1 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $4v_0 + v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| 3 3 2 2 1 | $2v_0 + 3v_1$ | $v_0 + 4v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $9v_0 + 16v_1$ |
| 3 2 2 3 1 | $v_0 + 2v_1$ | $v_0 + 2v_1$ | $v_0 + 2v_1$ | $v_0 + 2v_1$ | $3v_1$ | $4v_0 + 11v_1$ |
| 3 3 3 3 1 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| 3 2 3 2 2 | $2v_0 + 3v_1$ | $4v_0 + v_1$ | $2v_0 + 3v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $14v_0 + 11v_1$ |
| 3 3 3 3 3 | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $5v_0 + 20v_1$ |

Pareto equilibria when $\geq 1/2$, NB solutions 81, Nb distinct equilibria : 12

| Local functions | Players utility [$\times 6$] | | | | | $SW[\times 30]$ |
|-----------------|--------------------------------|---------------|---------------|---------------|---------------|-----------------|
| 2 1 1 0 0 | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $5v_0$ | $13v_0 + 12v_1$ |
| 3 2 1 1 0 | $2v_0 + 3v_1$ | $4v_0 + v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $11v_0 + 14v_1$ |
| 1 3 1 1 0 | $5v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $7v_0 + 18v_1$ |
| 1 3 3 1 0 | $5v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $5v_1$ | $5v_0$ | $7v_0 + 18v_1$ |
| 2 3 1 2 0 | $3v_0 + 2v_1$ | $v_0 + 4v_1$ | $5v_1$ | $3v_0 + 2v_1$ | $5v_0$ | $12v_0 + 13v_1$ |
| 3 2 2 3 0 | $v_0 + 4v_1$ | $4v_0 + v_1$ | $4v_0 + v_1$ | $v_0 + 4v_1$ | $5v_0$ | $15v_0 + 10v_1$ |
| 2 2 1 1 1 | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_1$ | $6v_0 + 19v_1$ |
| 3 3 1 2 1 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $4v_0 + v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| 3 3 2 2 1 | $2v_0 + 3v_1$ | $v_0 + 4v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $9v_0 + 16v_1$ |
| 3 3 3 3 1 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| 3 2 3 2 2 | $2v_0 + 3v_1$ | $4v_0 + v_1$ | $2v_0 + 3v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $14v_0 + 11v_1$ |
| 3 3 3 3 3 | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $v_0 + 4v_1$ | $5v_0 + 20v_1$ |

Nash equilibria for NC_{01}

Nash equilibria for NC_{01} when $1/3 \leq v_0/v_1 \leq 1/2$, Nb solutions : 76, Nb

distinct equilibria : 13

| Local functions | Players utility [$\times 6$] | | | | | $SW[\times 30]$ |
|-----------------|--------------------------------|---------------|---------------|---------------|---------------|-----------------|
| 1 1 2 0 0 | $5v_1$ | $5v_1$ | $2v_0 + 3v_1$ | $5v_0$ | $5v_0$ | $12v_0 + 13v_1$ |
| 3 2 1 1 0 | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $11v_0 + 14v_1$ |
| 1 3 1 1 0 | $5v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $8v_0 + 17v_1$ |
| 3 2 2 1 0 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $5v_0$ | $11v_0 + 14v_1$ |
| 1 3 3 1 0 | $5v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $5v_0$ | $9v_0 + 16v_1$ |
| 1 3 1 2 0 | $5v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $2v_0 + 3v_1$ | $5v_0$ | $9v_0 + 16v_1$ |
| 2 1 3 2 0 | $2v_0 + 3v_1$ | $5v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $5v_0$ | $11v_0 + 14v_1$ |
| 2 2 1 1 1 | $3v_0 + 2v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $5v_1$ | $5v_1$ | $5v_0 + 20v_1$ |
| 3 3 1 2 1 | $2v_0 + 3v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| 3 3 2 2 1 | $3v_0 + 2v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $10v_0 + 15v_1$ |
| 3 3 3 3 1 | $3v_0 + 2v_1$ | $2v_0 + 3v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $11v_0 + 14v_1$ |
| 3 2 3 2 2 | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $2v_0 + 3v_1$ | $14v_0 + 11v_1$ |
| 3 3 3 3 3 | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $2v_0 + 3v_1$ | $10v_0 + 15v_1$ |

Nash equilibria for NC_{01} when $v_0/v_1 \geq 1/2$, Nb solutions 40, Nb distinct equi-

libria: 6

| Local functions | Players utility [$\times 6$] | | | | | $SW[\times 30]$ |
|-----------------|--------------------------------|---------------|---------------|---------------|---------------|-----------------|
| 3 2 1 1 0 | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $11v_0 + 14v_1$ |
| 1 3 1 1 0 | $5v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $5v_1$ | $5v_0$ | $8v_0 + 17v_1$ |
| 2 2 1 1 1 | $3v_0 + 2v_1$ | $2v_0 + 3v_1$ | $5v_1$ | $5v_1$ | $5v_1$ | $5v_0 + 20v_1$ |
| 3 3 1 2 1 | $2v_0 + 3v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $8v_0 + 17v_1$ |
| 3 3 3 3 1 | $3v_0 + 2v_1$ | $2v_0 + 3v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $5v_1$ | $11v_0 + 14v_1$ |
| 3 2 3 2 2 | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $3v_0 + 2v_1$ | $2v_0 + 3v_1$ | $14v_0 + 11v_1$ |

References

- [1] Anurag Anshu, Peter Høyer, Mehdi Mhalla, and Simon Perdrix. Contextuality in multipartite pseudo-telepathy graph games. In *International Symposium on Fundamentals of Computation Theory*, pages 41–55. Springer, 2017.
- [2] Vincenzo Auletta, Diodato Ferraioli, Ashutosh Rai, Giannicola Scarpa, and Andreas Winter. Belief-invariant and quantum equilibria in games of incomplete information. *arXiv preprint arXiv:1605.07896*, 2016.
- [3] Robert J. Aumann. Subjectivity and correlation in randomized strategies. *Journal of mathematical Economics*, 1(1):67–96, 1974.
- [4] Manik Banik, Some Sankar Bhattacharya, Nirman Ganguly, Tamal Guha, Amit Mukherjee, Ashutosh Rai, and Arup Roy. Two-Qubit Pure Entanglement as Optimal Social Welfare Resource in Bayesian Game. *Quantum*, 3:185, Sep. 2019.
- [5] Katarzyna Bolonek-Lasoń. Three-player conflicting interest games and nonlocality. *Quantum Information Processing*, 16(8):186, 2017.
- [6] Gilles Brassard, Anne Broadbent, and Alain Tapp. Quantum pseudo-telepathy. *Foundations of Physics*, 35(11):1877–1907, 2005.
- [7] Harry Buhrman, Oded Regev, Giannicola Scarpa, and Ronald De Wolf. Near-optimal and explicit bell inequality violations. In *2011 IEEE 26th Annual Conference on Computational Complexity*, pages 157–166. IEEE, 2011.
- [8] Matthew Coudron, Jalex Stark, and Thomas Vidick. Trading locality for time: certifiable randomness from low-depth circuits. *arXiv preprint arXiv:1810.04233*, 2018.
- [9] Françoise Forges. Correlated equilibria in a class of repeated games with incomplete information. *International Journal of Game Theory*, 14:129–149, 1985.
- [10] Françoise Forges. Five legitimate definitions of correlated equilibrium in games with incomplete information. *Theory and decision*, 35(3):277–310, 1993.

- [11] Françoise Forges. Correlated equilibrium in games with incomplete information revisited. *Theory and decision*, 61(4):329–344, 2006.
- [12] Justin Holmgren and Lisa Yang. The parallel repetition of non-signaling games: Counterexamples and dichotomy. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2019, page 185–192, New York, NY, USA, 2019. Association for Computing Machinery.
- [13] Marius Junge, Carlos Palazuelos, David Pérez-García, Ignacio Villanueva, and Michael M Wolf. Unbounded violations of bipartite bell inequalities via operator space theory. *Communications in Mathematical Physics*, 300(3):715–739, 2010.
- [14] Ehud Lehrer, Dinah Rosenberg, and Eran Shmaya. Signaling and mediation in games with common interests. *Games and Economic Behavior*, 68(2):670–682, 2010.
- [15] Qingmin Liu. Correlation and common priors in games with incomplete information. *Journal of Economic Theory*, 157:49–75, 2015.
- [16] Louis Mathieu and Mehdi Mhalla. Separating pseudo-telepathy games and two-local theories. *arXiv preprint arXiv:1806.08661*, 2018.
- [17] John F Nash et al. Equilibrium points in n-person games. *Proceedings of the national academy of sciences*, 36(1):48–49, 1950.
- [18] Anna Pappa, Niraj Kumar, Thomas Lawson, Miklos Santha, Shengyu Zhang, Eleni Diamanti, and Iordanis Kerenidis. Nonlocality and conflicting interest games. *Physical Review Letters*, 114(2):020401, 2015.
- [19] Arup Roy, Amit Mukherjee, Tamal Guha, Sibasish Ghosh, Some Sankar Bhattacharya, and Manik Banik. Nonlocal correlations: Fair and unfair strategies in bayesian games. *Phys. Rev. A*, 94:032120, Sep. 2016.